

Computational Modelling Folate Metabolism and DNA Methylation: Implications for Understanding Health and Ageing

Mark T. Mc Auley¹, Kathleen M. Mooney² and J. Enrique Salcedo-Sora³

¹Department of Chemical Engineering, Thornton Science Park, University of Chester, CH2 4NU, UK

²Faculty of Health and Social Care, Edge Hill University, Ormskirk, Lancashire, L39 4QP, UK

³School of Health Science, Liverpool Hope University, Hope Park, Liverpool, L16 9JD, UK

Author Description

Mark Mc Auley is a Computational Systems Biologist and Senior Lecturer of Chemical Engineering in the Faculty of Science and Engineering, University of Chester, UK. His research interests are computational modelling of nutrient based systems and their interaction with ageing. Specifically, folate and lipid metabolism.

Kathleen Mooney is a Nutritionist who works as a Senior Lecturer in Nutrition at Edge Hill University, UK. Her interests are the role of B vitamins in maintaining health.

Enrique Salcedo-Sora has a Medical Degree and a Master of Science/Ph.D. in Biochemistry. Currently he works as a Lecturer in Medical Biochemistry in the School of Health Sciences at Liverpool Hope University, UK. He is also an Honorary Research Fellow at the Liverpool School of Tropical Medicine, UK. He is interested in folate metabolism in both eukaryotic and prokaryotic organisms and how perturbations to folate metabolism can be used to help treat diseases such as malaria.

Abstract

Dietary folates have a key role to play in health as deficiencies in the intake of these B vitamins have been implicated in a wide variety of clinical conditions. The reason for this is folates function as single carbon donors in the synthesis of methionine and nucleotides. Moreover, folates have a vital role to play in the epigenetics of mammalian cells by supplying methyl groups for DNA methylation reactions. Intriguingly, a growing body of experimental evidence suggests DNA methylation status could be a central modulator of the ageing process. This has important health implications because the methylation status of the human genome could be used to infer age-related disease risk. Thus, it is imperative we further our understanding of the processes which underpin DNA methylation and how these intersect with folate metabolism and ageing. The biochemical and molecular mechanisms which underpin these processes are complex. However, computational modelling offers an ideal framework for handling this complexity. A number of computational models have been assembled over the years, but to date no model has represented the full scope of the interaction between the folate cycle and the reactions which govern the DNA methylation cycle. In this review we will discuss several of the models which have been developed to represent these systems. In addition we will present a rationale for developing a combined model of folate metabolism and the DNA methylation cycle.

Keywords: folate metabolism, computational model, DNA methylation, ageing, stochastic [model](#)

Introduction

The term 'folate' (vitamin B9) is used to denote a group of compounds which possess the same vitamin activity and includes natural folates as well as the pharmacological compounds folic acid and folinic acid [1]. The dietary importance of folate cannot be overstated. This is emphasized by the wide range of clinical disorders which correlate with low folate status [2-4]. In addition, polymorphisms in genes coding for folate dependent enzymes [5, 6] are associated with several cancers (cervical, bronchial, colon and breast) [7-10], Alzheimer's disease [11], Down syndrome [12], unexplained recurrent early pregnancy loss and pre-eclampsia [13]. These associations are unsurprising when one considers that folates are involved in a ubiquitous array of cellular processes. For instance, folates are involved in the synthesis of nucleotides from purine precursors, participate indirectly in the synthesis of transfer RNA and function as single carbon donors during the re-methylation of homocysteine (Hcy) to methionine [14] (Figure 1). Folate-derived one-carbon units also play a central role in DNA methylation [15, 16]. This epigenetic process involves methyl groups covalently bonding to CpG dinucleotides to establish tissue specific methylation patterns. A CpG dinucleotide consists of a deoxycytidine followed by a deoxyguanine, with the "p" indicating the phosphate group between these nucleotides. Covalent bonding of methyl groups occurs at the carbon-5 position of a deoxycytidines to create a methylated CpG dyad [11, 12]. Dynamic changes to methylation patterns are an important gestational phenomenon, which

regulate gene expression during embryonic development [17]. However, during adult life alterations to DNA methylation patterns can have significant implications for the onset of disease [18, 19]. Specifically, advancing age is often accompanied by global hypomethylation [20-23] in conjunction with site specific hypermethylation at the promoter region of a variety of genes [24-27]. Hypermethylation, involves CpG islands (~500-2000 base pairs) within the promoter becoming excessively methylated [28, 29], and often occurs with concomitant transcriptional silencing [30, 31]. Gene promoter hypermethylation is a feature of certain diseases, the most notable being cancer [25]. [For instance, Wang et al. \(2016\) \[32\] recently found that DNA methylation changes during ageing were closely correlated to the occurrence of cancer.](#) In addition to cancer aberrant DNA methylation has an emerging role to play in many age related diseases including, cardiovascular disease (CVD) [33], Alzheimer's disease [34] and osteoporosis/osteoarthritis [35]. Such findings consolidate the growing view that DNA methylation status and health-span are inexorably entwined. Moreover it has also been revealed that investigating changes to the methylation profile of the genome could help develop our understanding of ageing. This assertion is supported by the recent bioinformatics work of [Horvath \(2013\) \[36\]](#) who utilized publicly available methylation data sets to identify an "epigenetic clock " underpinned by methylation changes in 353 CpGs. Intriguingly, Horvath postulates that "DNA methylation age" measures the cumulative effect of an epigenetic maintenance system. This finding indicates that a deeper mechanistic understanding of DNA methylation preservation could be pivotal to improving our overall understanding of ageing and health-span.

To extend our understanding of DNA methylation and its bidirectional relationship with ageing it is imperative we consider how subtle alterations to folate one

carbon metabolism (FOCM) impact the regulatory [processes](#) which govern DNA methylation. Specifically, it is necessary to explore how such perturbations [affect](#) gene promoters which are susceptible to hypermethylation. In addition, it is vital we consider how other factors associated with ageing impact this relationship. However, this is not a straightforward task as both FOCM and the DNA methylation cycle are complex processes which are underpinned by a large number of biochemical and molecular reactions (Figure 1). Many of these reactions are non-linear in nature and are influenced by a variety of other B vitamins/nutrients and enzymatic cofactors [37]. Ageing adds a further degree of complexity by altering the dynamics of the reactions. For example, age-dependent decreases in the expression levels/activity of human [methionine synthase \(MS\)](#) and methylenetetrahydrofolate reductase (MTHFR) have been observed [38, 39]. Both enzymes are recognised as key regulators of FOCM and expression/activity changes potentially affect their reaction kinetics. Similarly, the activity of human DNA methyltransferase 1 (Dnmt 1), a key enzyme involved in the addition of methyl groups to cytosine residues, has also been observed to decrease with age [40]. Moreover, ageing also affects the availability of cofactors such as vitamin B12 [41], which is central to FOCM, while oxidative stress, which is generally regarded as a key contributor to ageing, has been observed to effect both DNA methylation and FOCM [42, 43]. Furthermore, it is not uncommon that older people have diminished folate status as a result of a low dietary intake of the vitamin [44]. Thus, it is necessary to consider an array of factors when investigating the maintenance of DNA methylation. Fortunately, there is a growing appreciation that complex biological process can be studied in a holistic manner by adopting a systems biology approach [45]. Computational modelling resides at the centre of this paradigm shift as it provides a framework for representing and exploring the dynamics of complex systems [46-50]. In this

review we discuss the role computational modelling has played in developing our understanding of FOCM and DNA methylation. In addition we propose coupling FOCM and the DNA methylation cycle into one computational model, which could be used to further explore the dynamics of their relationship.

Folate metabolism and the DNA methylation Cycle

FOCM is fundamentally important to DNA methylation. Various folates are the cofactors for the *de novo* synthesis of methyl groups from more oxidized one carbon units and 2) the methyl groups on 5-methyltetrahydrofolate are used to re-methylate hcy to methionine and to regenerate tetrahydrofolate, the metabolically active form of folate. Methionine is a precursor of S-adenosylmethionine (SAM), which has a predominant role to play in the majority of biochemical methyl donation events, including that of DNA methylation. Post replication, Dnmt1 uses SAM as a substrate to transfer methyl groups to the DNA molecule [51] this produces S-adenosylhomocysteine (SAH). SAH is then converted to Hcy permitting the continuation of the methylation cycle [52, 53] (Figure 1). In conjunction with Dnmt1 DNA methylation patterns are dynamically regulated by several other enzymes (Figure 1). As, Dnmt1 preferentially acts on hemimethylated DNA and is potentially unable to methylate neo-synthesized DNA strands, this enzyme is generally regarded as solely a maintenance enzyme [54]. Thus, other enzymes are needed to perform *de novo* DNA methylation. Dnmt3a and Dnmt3b are widely regarded as the main enzymes responsible for this role [55, 56]. These methylation reactions are offset by active and passive demethylation. Passive demethylation occurs during replication, while active demethylation can involve ten eleven translocation (TET) dioxygenases, which oxidize the methyl groups of cytosine [57]. TET are thought to function by oxidizing the methyl groups of cytosine. This process eventually culminates with the reintroduction of unmethylated cytosine in the DNA molecule [57]. Thus, it

would appear that the steady state levels of both site specific and global DNA methylation patterns are maintained by an antagonistic balancing act between those activities responsible for maintenance/*de novo* methylation and those reactions responsible for passive/active demethylation. Moreover, methylation fidelity studies suggest these processes are subject to inherent stochasticity. For example, [Landan et al. \(2012\) \[58\]](#) tracked the *in vitro* evolution of immortalized fibroblasts for >300 generations and found that changes in population-averaged methylation occur through a stochastic process. In addition, [Shipony et al. \(2014\) \[59\]](#) suggest the persistence nature of the somatic methylome is one factor which makes it increasingly vulnerable to noise with time. This is intriguing from the perspective of ageing [as it would appear](#) that in young somatic cells, site specific methylation density is characterised by low level noise, which maintains average methylation density. Therefore, the persistent nature of the methylome and increased stochasticity with time could contribute to the formation of aberrant DNA methylation patterns which are a hallmark of the ageing process in humans.

Computational Models of Folate Metabolism

[The models described in the next two sections are summarised in Table 1 and 2 respectively.](#) Models of FOCM have traditionally adopted a continuous deterministic approach, with the dynamics of the system being represented by ordinary differential equations (ODEs), [which are underpinned by kinetic reactions \[60\].](#) [In the main reaction velocities are assumed to depend on the concentrations of the reaction substrates \(the law of mass action \[61\]\).](#) Rate laws for more complex reactions laws are in general described by Michaelis–Menten kinetics [62] for one, two, or three substrates assuming a random-order mechanism. The following mathematical expressions taken from the recent model by [Salcedo-Sora and Mc Auley \(2016\) \[63\]](#) encapsulate the different

Michaelis–Menten based functional forms as used for one , two, and three substrates in these types of kinetic models, where V is the velocity of the reaction, V_{\max} , is the theoretical maximum velocity of the reaction, S_1 to S_3 are substrates and k_{m1} to k_{m3} are the substrate concentrations at which the reaction is half maximum velocity.

$$V = \frac{V_{\max} * [S]}{k_m + [S]} \quad [1]$$

$$V = \frac{V_{\max} * [S_1][S_2]}{k_{m_1}k_{m_2} + k_{m_1}[S_2] + k_{m_2}[S_1]} \quad [2]$$

$$V = \frac{V_{\max} * [S_1][S_2][S_3]}{k_{m_1}k_{m_2}k_{m_3} + k_{m_1}([S_2] + [S_3]) + k_{m_2}([S_1] + [S_3]) + k_{m_3}([S_1] + [S_2]) + [S_1][S_2][S_3]} \quad [3]$$

The first mathematical model to adopt this kinetic approach to represent intracellular folate metabolism was developed in the 1970s and had a strong pharmacological theme [64]. It was used to simulate the actions of methotrexate on DNA synthesis. The next significant computational model of the folate cycle, was developed by [Nijhout *et al.* \(2004\) \[65\]](#). This model is underpinned by the enzyme kinetic data that characterises the reactions in Figure 1, [and was assembled using the type of mathematical equations outlined above](#). As well as being the first detailed mathematical description of the folate cycle, model simulations were able to quantitatively reproduce the intracellular levels of the various folate metabolites and the model was able to predict the effect of vitamin B12 deficiency. Building on this work, [Reed *et al.* \(2006\) \[66\]](#) used this model combined with models of methionine metabolism [67, 68] to investigate genetic polymorphisms in the folate pathway. Polymorphisms in MTHFR predicted a reduction in MTHFR activity reduces concentrations of SAM and 5-MTHR, and

DNA methylation, while slightly increasing SAH and hcy concentrations and thymidine or purine synthesis. Decreased folate together with a simulated vitamin B12 deficiency resulted in decreased DNA methylation and purine and thymidine synthesis. This model was also used as a template to study the effect of intracellular folate deficiency and excess. The model suggested that the enzyme thymidylate synthesis is very sensitive to changes in epithelial intracellular folate and increased significantly under conditions of high intracellular folate [69]. This framework was further extended by [Duncan et al. \(2013\) \[70\]](#) to hepatic and plasma folate turnover. It was applied to a population of virtual individuals and showed that tissue and plasma folate is highly correlated, but liver and plasma folate much less so. Moreover, this model showed that oxidative stress increases the plasma S-adenosylmethionine/S-SAM/SAH ratio. The most recent ODE model of folate metabolism was developed by [Salcedo-Sora and Mc Auley \(2016\) \[63\]](#). Our model is the first computational model of microbial folate biosynthesis and utilisation. Using the model we were able to identify specific targets within folate metabolism which synergise with current antifolates. In addition the results from our model support experimental findings which suggest the folinic acid substrate cycle is an important biochemical mechanism deployed during active cell growth. A key finding which emphasizes the utility of computational models for shedding light on this complex system.

Computational Models of the DNA Methylation Cycle

A variety of approaches have been used to model various aspects of DNA methylation [71-79]. Our focus is the precise representation of the DNA methylation cycle. To this end the DNA methylation cycle has been modelled explicitly by McGovern et al. (2012) [80]. The model centred on the activity of DNMTs and partial differential equations [was](#) used to represent methylation

reactions. It was used as a predictive tool for haematological malignancies. The model was able to predict the relative abundances of unmethylated, hemimethylated, fully methylated, and hydroxymethylated CpG dyads in the DNA of cells with fully functional methylation and TET enzymes. In addition to this deterministic framework stochastic models of the methylation cycle have also been used to study the methylation cycle. The goal has been to derive models which account for the noise associated with maintaining DNA methylation levels. For example, a reduced probabilistic mathematical representation of methylation dynamics (known as the standard/classical model) has been proposed by Pfeifer *et al.* (1990) [81], and further utilised by Riggs and Xiong (2004) [82]. This model is presented in equations 4-7.

$$\frac{dM}{dt} = E_m * M + E_d * U \quad [4]$$

$$\frac{dU}{dt} = (1 - E_d) * U + (1 - E_m) * M \quad [5]$$

$$M^{\diamond} = \frac{M}{Total} = \frac{M}{(M + U)} \quad [6]$$

$$M^{\diamond} = \frac{E_d}{(1 + E_d - E_m)} \quad [7]$$

To briefly explain ODEs 4 and 5, and equations 6 and 7, M is a CpG site, U is its unmethylated state, E_d is the efficiency of *de novo* methylation, E_m is the efficiency of maintenance methylation, M is the number of methylated sites, M^{\wedge} is the fraction of methylation, while equation 7 represents the steady state solution to ODEs 4 and 5 when $dM^{\wedge}/dt=0$. This model is underpinned by the idea that site specific methylation levels are underscored by a stochastic process, where *de novo* and maintenance methylation activity are central. Recently Jeltsch and Jurkowska (2014) [83] provided an intriguing theoretical extension to this concept. Jeltsch and Jurkowska propose a stochastic model which also includes the rate of cell division together with passive and active demethylation

(equation 8). θ_m^i , is the change in the fraction of methylation at any CpG site (i). This is determined by the difference between the rate of *de novo* DNA methylation (r_{met}^i) and rates of loss of methylation by replication and active demethylation (r_{demeth}^i) at this site. D , is the rate of cell division, f_{main}^i is the efficiency of maintenance methylation at site i.

$$\frac{d\theta_m^i}{dt} = r_{met}^i * (1 - \theta_m^i) - \left(\frac{1}{2} D (1 - f_{main}^i) + r_{demeth}^i \right) * \theta_m^i \quad [8]$$

According to the authors, equation 8 has the significant advantage over its predecessor as it includes the possibility of proofreading/repair mechanisms due to the incorporation of negative feedback. This level of regulation is attained because an increase in methylation causes a reduction in the first term of equation 8 with a subsequent increase in the second term. Consequently the methylation rate drops and demethylation increases, resulting in a stable steady-state methylation level at each site. To our knowledge this theoretical framework has yet to be applied to experimental data or simulated dynamically. A stochastic mathematical framework also inspired Haerter *et al.* (2014) [84] to create a computational model of this system. Their model differs from the classical/standard model as it included dynamic collaboration between CpGs. More recently Olariu *et al.* (2016) [85] built on the work of Haerter *et al.* (2014) to model the regulation of transcription and DNA methylation.

Rationale for Computationally Integrating FOCM and the DNA Methylation Cycle

Computational models of FOCM are not fully integrated with the DNA methylation cycle. Rather, as outlined they have been restricted to the cycle itself or its

involvement with DNA synthesis and have tended not to include the DNA methylation cycle. Some have attempted to account for DNA methylation but no model to our knowledge has explicitly integrated both systems in a meaningful way. We propose computationally integrating FOCM and the methylation cycle. Our biological rationale is unpinned by the knowledge that perturbations in the folate cycle are strongly coupled with aberrant DNA methylation. This is evidenced by the association between low folate status, global DNA hypomethylation and increased cancer risk [86]. In addition, genetic polymorphisms most notably in the MTHFR gene, which codes for [MTHFR](#), an enzyme which catalyses the conversion of 5, 10-methylenetetrahydrofolate to 5-methyltetrahydrofolate (5-MTHF) (Figure 1) is associated with gene promoter hypermethylation [87]. Moreover, methionine synthase A2756G (MTR A2756G) is a common polymorphism in the gene encoding [MS](#). This enzyme catalyses the regeneration of methionine from hcy and disruptions to its enzymatic capabilities also emphasise the close link between folate metabolism and disruptions to DNA methylation. For instance, [Weiner and colleagues \(2014\)](#) [87] found that individuals homozygous for this mutant allele showed higher leukocyte genomic DNA methylation levels than individuals possessing the wild-type genotype (MTR 2756AA). It makes mechanistic sense that perturbations of this nature to FOCM have an impact on DNA methylation, as 5-methyltetrahydrofolate, is a cosubstrate for hcy remethylation to methionine, while MS modulates the dynamics of this reaction and perturbations to this enzyme result in a rise in plasma hcy [88], which is associated with CVD [89]. Intriguingly, a rise in plasma hcy has also been associated with gene promoter hypermethylation [90-93] and global hypomethylation [94-96] of human DNA.

In addition to perturbations due to genetic mutations it is important to recognise that FOCM compartmentalisation could have implications for DNA methylation status as compartmentalisation of FOCM has a role to play in regulating the distribution of one-carbon units within the cell. The partitioning of one-carbon units between nucleotide synthesis and hcy remethylation is the focal point of this process, where methylenetetrahydrofolate (MTHF) is at the centre of a metabolic competition [97, 98]. On the one hand MTHF is used as a cofactor during nucleotide synthesis to facilitate the conversion of dUMP to dTMP by thymidylate synthase (TYMS). However, MTHF can also be reduced and is then fully committed to the remethylation of hcy. The separation of FOCM into distinct cellular compartments is the determining factor which regulates the competition between TYMS and MTHFR for MTHF [99]. This regulation requires the trifunctional, enzyme methylenetetrahydrofolate dehydrogenase (MTHFD1), whose activity depends on its cellular location [99]. The interdependence of this enzymes activity to its location and its involvement in regulating the competition between MTHFR and TYMS is evident, when intracellular folate levels are deficient, as insufficient levels of folate cause an accumulation of MTHFD1 in the nucleus, when compared to its cytosolic counterpart [100]. The nuclear build-up of MTHFD1 emphasises FOCM will sacrifice hcy remethylation in favour of nucleotide synthesis and underscores the regulatory role compartmentalisation plays in one-carbon metabolism. Intriguingly this aspect of FOCM could have wider implications as recently it has been shown that changes to MTHFD1 can affect the regulation of DNA methylation. Groth *et al.* (2016) [101] identified a mutant (*mthfd1-1*) in *Arabidopsis thaliana*, which carried a mutation in cytoplasmic MTHFD1. This mutant suffered from accumulation of hcy and SAM, coupled with extensive genome-wide hypomethylation. Compartmental regulation of FOCM is also highlighted by cyserine hydroxymethyltransferase

(SHMT). SHMT1 catalyses the conversion of tetrahydrofolate (THF) to 5, 10-methylenetetrahydrofolate and also converts serine to glycine. It has been found that cytosine serine hydroxymethyltransferase (cSHMT1) preferentially supplies one-carbon units for thymidylate synthesis [98, 102]. Mechanistically, it has been found this preference is mediated by small ubiquitin-like modifier (SUMO) modification. This alteration enables the translocation of SHMT to the nucleus and also modifies its catalytic capabilities [103]. Sumoylation could also be crucial to both DNA methylation and demethylation as SUMO-1 modification of DNMT1 had been shown to significantly enhance DNMT1 activity [104]. It is also possible sumoylation is capable of modulating the activity of Dnmt3a and Dnmt3b [105]. This level of crossover between FOCM and DNA methylation consolidates the view that disruptions to FOCM impact the dynamics of DNA methylation/demethylation events. However, the full extent of this relationship remains to be elucidated, hence the growing need for an integrated computational model to explore the complex interactions of these systems.

Theoretical Frameworks and Obstacles to Creating a Combined Model

To develop a combined model several theoretical approaches could be adopted. FOCM could be retained as a set of kinetic based ODEs, as is the case in the majority of models of this nature. It would then be straightforward to add the methylation cycle as ODEs also and to define the combined system as a coupled set of ODEs. Despite the obvious computational advantages of this simplification, these benefits are offset by several factors. Firstly, to assemble this deterministic model the ODEs need to be informed by kinetic data, which accounts for the behaviour of the methylation enzymes. [This is not a straightforward task as parameter uncertainty is a significant issue in the field of computational systems](#)

biology. As outlined, kinetic based models of folate metabolism tend to have rate laws where the velocity of a reaction assumes mass action kinetics or is based on an enzyme kinetic law (e.g. Michaelis-Menten kinetics). Unfortunately kinetic parameters can vary significantly depending on the circumstances in which their kinetics were quantified and the biological source, with the possibility of model parameters varying by several orders of magnitude depending on the particular source [106]. Therefore, it is extremely unlikely that a mathematical model can be assembled from one biological source. This is highlighted by our recent model of microbial folate metabolism, where the parameters were derived from a number of sources [63]. It is also a feature of the mammalian computational models which have previously relied on data from rat, mice and human studies [65]. Despite such pitfalls, there is no alternative to this approach of determining model parameters. To this end, a number of databases now exist which archive kinetic data for the purpose of assembling computational models [107, 108]. We used these databases to identify a series of indicative kinetic parameters for the model outlined diagrammatically in Figure 1 and are summarised in Table 3. The identification of these parameters serve to emphasize that it is possible to assemble a combined kinetic model of folate metabolism and the DNA methylation cycle based on human data, however the data is limited. Moreover, we were able to identify kinetic parameters for Dnmt1 and Dnmt3A [109, 110], however kinetic information for the TET demethylation enzymes could not be identified.

Although as outlined above it is clearly possible to assemble a combined deterministic kinetic model, given experimental evidence indicates DNA methylation events are underscored by stochasticity a kinetic based deterministic approach is not the correct way to model this aspect of the system.

A more accurate way is to incorporate noise into the model. The standard way of representing noise within biological cells is to describe it mathematically by a master equation or to simulate molecule fluctuations by using a stochastic simulation algorithm (SSA), such as the Gillespie algorithm or one of its derivatives [111, 112]. Routinely embedded within the idea of intracellular noise is the concept of intrinsic versus extrinsic noise, as developed experimentally by Elowitz and colleagues (2002) [113]. Within this context intrinsic noise refers to the variability inherent to the system under consideration, while fluctuations in those factors classified as external to the system of interest are responsible for extrinsic noise (Figure 2). Specifically, this mathematical approach has been ubiquitously applied to the study of cell to cell variability in gene expression levels within isogenic cell populations [114-117].

Applying the above logic to our indicative model it can be argued the reactions of the methylation cycle are responsible for generating intrinsic noise e.g. the inherent statistical mechanical fluctuations in the binding and diffusion dynamics of the molecules involved in maintaining methylation levels within a CpG island/[gene body](#). Conversely, DNA methylation/the epigenetic state of the cell is considered to be a component of extrinsic noise when variations in gene expression are modelled in this way. We suggest extrinsic noise in our integrated model arises due to biochemical fluctuations originating from the folate cycle and other cell parameters particularly those related to ageing (Figure 2). Including biochemical noise arising from FOCM is crucial especially if one considers that site specific (e.g. gene promoter) DNA methylation events interact with FOCM within a microscopic rather than macroscopic environment. This microenvironment is characterised by low molecular populations, which react at discrete time-points, via random collisions between individual molecules. The

advantage of this framework is its simplicity as a complex biological network is reduced to a much simpler abstraction. A drawback of this approach is that it reduces the methylation cycle and FOCM to two variables within a phenomenological model, which lumps all sources of extrinsic noise together. Moreover, as we are concerned with the complexities of the reactions of folate metabolism and how changes to these impact DNA methylation events, it is challenging to separate this source of noise from other extrinsic events e.g. the impact of the cell cycle. A further challenge associated with this proposal is that in order to characterise the intrinsic noise for a given gene promoter it would be necessary to experimentally characterise promoter specific cell to cell methylation variability in young disease free cells and how this relates to protein expression levels. For instance, microscopy or flow cytometry are routinely used to quantify protein expression variability in reporter genes embedded in homogenous cell populations [118, 119]. Protein count distributions then provide an overview of cell to cell fluctuations in gene expression. The stochastic profile can then be characterised as the squared coefficient of variation of the fluorescence levels [120], the coefficient of variation [113] or the Fano Factor [121]. Therefore, it would be necessary to develop experimental methods which are capable of determining the exact nature of the stochasticity of the DNA methylation process. However, given that DNA methylation levels are quantified in a totally different manner to protein fluctuations it is could be challenging to define/characterise its stochastic signature in a precise way experimentally. Next generation sequencing techniques could help to alleviate this problem. These techniques enable the determination of the methylation status at each cytosine for a specific segment of the genome. An eloquent example of this approach is the work of Hansen et al. (2011) [122], who showed stochastic methylation of cancer-specific DNA-methylated regions. This was able to distinguish cancer from

normal tissue. More recently Cheow *et al.* (2016) [123] developed a method to genotype individual cells while concurrently probing gene expression and DNA methylation levels at multiple loci. Such methods could help identify an individual stochastic signature for a particular gene promoter/gene body, which would help inform the mathematics underpinning future computational models. An alternative to the reduced stochastic model approach would be to explicitly represent each deterministic kinetic reaction within FOCM and the DNA methylation cycle as stochastic propensity functions and to simulate the integrated model with a SSA [99], as has been done with other complex biochemical systems [124]. In Figure 2 part b we have created a list of reactions which could be represented as propensity functions, where C_1 - C_7 are indicative of stochastic rate constants. This is not unrealistic as mathematical approaches have been developed previously for converting deterministic systems biology models into stochastic models [125-127]. However, regardless of the approach it is our opinion that future computational models should combine these two systems together. Moreover, it is imperative that any combined model accounts for the biochemical and molecular variability which is inherent to both systems. This will improve our overall understanding of how those gene promoters which appear to be susceptible to hypermethylation interact with both folate metabolism and the ageing process.

Conclusions

Folate metabolism and the DNA methylation cycle are inexorably entwined as the folate cycle provides the methyl groups used in DNA methylation reactions. Experimental evidence suggests the dynamics of the folate cycle and the DNA methylation cycle are subject to inherent stochasticity. Investigating this behaviour is challenging, however computational modelling offers an ideal framework for exploring the interactions between these two systems. In this

review we proposed building a computational model of folate metabolism coupled with the reactions of the methylation cycle. This model would help to further explore the dynamics of this relationship and could be used to investigate how disruptions to these processes result in aberrant DNA methylation status, specifically gene promoter hypermethylation.

Key Points

- Folate metabolism has a vital role to play in health and possibly intrinsic ageing
- Foliates supply methyl groups for DNA methylation reactions which are central to gene regulation.
- Dynamic computational models have been successfully used to study the complexities of folate metabolism and the DNA methylation cycle respectively, but no model has fully coupled both systems.
- To combine both systems computationally it is important to account for the inherent stochasticity which influences the dynamics of both systems.
- Incorporating inherent biochemical and molecular variability will provide an improved understanding of how perturbations to these systems impact the onset of disease and will help to further elucidate the bidirectional relationship between ageing and DNA methylation.

Glossary

Deterministic model

A model which assumes variability does not impact the system of interest. The model will produce the same output given the same initial conditions and parameters.

Extrinsic noise

Fluctuations in the factors external to the biological system of interest e.g. biochemical fluctuations in the folate pathway and its impact on gene expression.

Gillespie algorithm networks.	An algorithm used to generate stochastic models of reaction networks.
Intrinsic noise	Noise associated with a biological system of interest e.g. in the case of this review, site specific DNA methylation.
Stochastic model	A model grounded in probability theory which is used to represent the variability inherent in biological processes. In this review we suggest FOCM is combined with the DNA methylation cycle into a single stochastic model, to capture the inherent variability in body systems.

Figure 1. Folate metabolism and its intersection with DNA Methylation: Ingested folates, are the cofactors for the *de novo* synthesis of methyl groups. Methyl groups on 5-methyltetrahydrofolate are used to re-methylate homocysteine (Hcy) to methionine, with the aid of vitamin B12 and methionine synthase. This reaction regenerates tetrahydrofolate, the metabolically active form of folate. Methionine is a precursor of S-adenosylmethionine (SAM), which has a predominant role to play in the majority of biochemical methyl donation events, including that of DNA methylation. DNA methyltransferase 1 is central to maintaining genomic methylation patterns, while DNA methyltransferase 3a and 3b are both involved in *de novo* methylation reactions. Ten eleven translocation (TET) dioxygenases are actively involved in demethylating cytosine, while CH₃ groups are also lost passively. Pink circles indicate methylated CpG's.

Figure 2. Stochastic contributions to CpG island methylation: Methylation density is affected by intrinsic and extrinsic noise. On the one hand noise is necessary and evolution has finely tuned this system so that stochasticity helps to maintain average methylation density in young cells. However, the ageing process results in gradual increase in stochasticity which eventually culminates with promoter hypermethylation. During aging gene promoter methylation has been identified within a wide variety of genes involved in health-span/ageing (Refer to table 4 for full function and description). Included also are the equations which define intrinsic and extrinsic noise, which are discussed in the main text (Mathematical equations based on framework introduced by Elowitz colleagues (2002) [113]). Part B presents a summary of the combined model represented by a set of indicative reactions which could be modelled using propensity functions. Abbreviations: F, folate; THF, tetrahydrofolate; 5MTHF, 5-methyltetrahydrofolate; 5,10MTHF, 5,10 methylenetetrahydrofolate; Met, methionine; SAM, S-adenosylmethionine; SAH, S-adenosylhomocysteine; Hcy, homocysteine; DHF, dihydrofolate; 10Form, 10,formyl THF.

Tables 1 to 4.

References

1. Hoffbrand AV, Weir DG. The history of folic acid, *Br J Haematol* 2001;113:579-589.
2. Robinson K, Arheart K, Refsum H et al. Low circulating folate and vitamin B6 concentrations: risk factors for stroke, peripheral vascular disease, and coronary artery disease. European COMAC Group, *Circulation* 1998;97:437-443.
3. Ramos MI, Allen LH, Mungas DM et al. Low folate status is associated with impaired cognitive function and dementia in the Sacramento Area Latino Study on Aging, *Am J Clin Nutr* 2005;82:1346-1352.
4. Yakut M, Ustun Y, Kabacam G et al. Serum vitamin B12 and folate status in patients with inflammatory bowel diseases, *Eur J Intern Med* 2010;21:320-323.
5. De Bree A, Verschuren WM, Kromhout D et al. Homocysteine determinants and the evidence to what extent homocysteine determines the risk of coronary heart disease, *Pharmacol Rev* 2002;54:599-618.
6. Molloy AM. Genetic aspects of folate metabolism, *Subcell Biochem* 2012;56:105-130.
7. Butterworth CE, Jr. Folate status, women's health, pregnancy outcome, and cancer, *J Am Coll Nutr* 1993;12:438-441.
8. Piyathilake CJ, Johanning GL, Macaluso M et al. Localized folate and vitamin B-12 deficiency in squamous cell lung cancer is associated with global DNA hypomethylation, *Nutr Cancer* 2000;37:99-107.
9. Powers HJ. Interaction among folate, riboflavin, genotype, and cancer, with reference to colorectal and cervical cancer, *J Nutr* 2005;135:2960s-2966s.
10. Chen P, Li C, Li X et al. Higher dietary folate intake reduces the breast cancer risk: a systematic review and meta-analysis, *Br J Cancer* 2014;110:2327-2338.
11. Corrada MM, Kawas CH, Hallfrisch J et al. Reduced risk of Alzheimer's disease with high folate intake: the Baltimore Longitudinal Study of Aging, *Alzheimers Dement* 2005;1:11-18.
12. Patterson D. Folate metabolism and the risk of Down syndrome, *Downs Syndr Res Pract* 2008;12:93-97.
13. Cao Y, Zhang Z, Zheng Y et al. The association of idiopathic recurrent early pregnancy loss with polymorphisms in folic acid metabolism-related genes, *Genes Nutr* 2014;9:402.
14. Kelly GS. Foliates: supplemental forms and therapeutic applications, *Altern Med Rev* 1998;3:208-220.
15. Crider KS, Yang TP, Berry RJ et al. Folate and DNA methylation: a review of molecular mechanisms and the evidence for folate's role, *Adv Nutr* 2012;3:21-38.
16. Li E, Zhang Y. DNA methylation in mammals, *Cold Spring Harb Perspect Biol* 2014;6:a019133.
17. Messerschmidt DM, Knowles BB, Solter D. DNA methylation dynamics during epigenetic reprogramming in the germline and preimplantation embryos, *Genes Dev* 2014;28:812-828.
18. Feil R. Epigenetics, an emerging discipline with broad implications, *C R Biol* 2008;331:837-843.
19. Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals, *Nat Genet* 2003;33 Suppl:245-254.
20. Richardson BC. Role of DNA methylation in the regulation of cell function: autoimmunity, aging and cancer, *J Nutr* 2002;132:2401S-2405S.
21. Wilson VL, Smith RA, Ma S et al. Genomic 5-methyldeoxycytidine decreases with age, *J Biol Chem* 1987;262:9948-9951.

22. Fuke C, Shimabukuro M, Petronis A et al. Age related changes in 5-methylcytosine content in human peripheral leukocytes and placentas: an HPLC-based study, *Ann Hum Genet* 2004;68:196-204.
23. Wilson VL, Jones PA. DNA methylation decreases in aging but not in immortal cells, *Science* 1983;220:1055-1057.
24. Issa JP, Vertino PM, Boehm CD et al. Switch from monoallelic to biallelic human IGF2 promoter methylation during aging and carcinogenesis, *Proc Natl Acad Sci U S A* 1996;93:11757-11762.
25. Esteller M, Corn PG, Baylin SB et al. A gene hypermethylation profile of human cancer, *Cancer Res* 2001;61:3225-3229.
26. Maegawa S, Hinkal G, Kim HS et al. Widespread and tissue specific age-related DNA methylation changes in mice, *Genome Res* 2010;20:332-340.
27. Issa JP, Ottaviano YL, Celano P et al. Methylation of the oestrogen receptor CpG island links ageing and neoplasia in human colon, *Nat Genet* 1994;7:536-540.
28. Dong SM, Kim HS, Rha SH et al. Promoter hypermethylation of multiple genes in carcinoma of the uterine cervix, *Clin Cancer Res* 2001;7:1982-1986.
29. Harden SV, Tokumaru Y, Westra WH et al. Gene promoter hypermethylation in tumors and lymph nodes of stage I lung cancer patients, *Clin Cancer Res* 2003;9:1370-1375.
30. Kane MF, Loda M, Gaida GM et al. Methylation of the hMLH1 promoter correlates with lack of expression of hMLH1 in sporadic colon tumors and mismatch repair-defective human tumor cell lines, *Cancer Res* 1997;57:808-811.
31. Esteller M, Silva JM, Dominguez G et al. Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors, *J Natl Cancer Inst* 2000;92:564-569.
32. Wang Y, Zhang J, Xiao X et al. The identification of age-associated cancer markers by an integrative analysis of dynamic DNA methylation changes, *Sci Rep* 2016;6:22722.
33. Abi Khalil C. The emerging role of epigenetics in cardiovascular disease, *Ther Adv Chronic Dis* 2014;5:178-187.
34. De Jager PL, Srivastava G, Lunnon K et al. Alzheimer's disease: early alterations in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci, *Nat Neurosci* 2014;17:1156-1163.
35. Delgado-Calle J, Fernandez AF, Sainz J et al. Genome-wide profiling of bone reveals differentially methylated regions in osteoporosis and osteoarthritis, *Arthritis Rheum* 2013;65:197-205.
36. Horvath S. DNA methylation age of human tissues and cell types, *Genome Biol* 2013;14:R115.
37. Ulrey CL, Liu L, Andrews LG et al. The impact of metabolism on DNA methylation, *Human Molecular Genetics* 2005;14:R139-R147.
38. Muratore CR, Hodgson NW, Trivedi MS et al. Age-dependent decrease and alternative splicing of methionine synthase mRNA in human cerebral cortex and an accelerated decrease in autism, *PLoS One* 2013;8:e56927.
39. Spotila LD, Jacques PF, Berger PB et al. Age dependence of the influence of methylenetetrahydrofolate reductase genotype on plasma homocysteine level, *Am J Epidemiol* 2003;158:871-877.
40. Li Y, Liu Y, Strickland FM et al. Age-dependent decreases in DNA methyltransferase levels and low transmethylation micronutrient levels synergize to promote overexpression of genes implicated in autoimmunity and acute coronary syndromes, *Exp Gerontol* 2010;45:312-322.
41. Zhang Y, Hodgson NW, Trivedi MS et al. Decreased Brain Levels of Vitamin B12 in Aging, Autism and Schizophrenia, *PLoS One* 2016;11:e0146797.

42. Franco R, Schoneveld O, Georgakilas AG et al. Oxidative stress, DNA methylation and carcinogenesis, *Cancer Lett* 2008;266:6-11.
43. Hondorp ER, Matthews RG. Oxidative stress inactivates cobalamin-independent methionine synthase (MetE) in *Escherichia coli*, *PLoS Biol* 2004;2:e336.
44. Haller J. The vitamin status and its adequacy in the elderly: an international overview, *Int J Vitam Nutr Res* 1999;69:160-168.
45. Mooney KM, Morgan AE, Mc Auley MT. Aging and computational systems biology, *Wiley Interdiscip Rev Syst Biol Med* 2016;8:123-139.
46. Mc Auley MT, Choi H, Mooney K et al. Systems Biology and Synthetic Biology: A New Epoch for Toxicology Research, *Advances in Toxicology* 2015;2015:14.
47. Mc Auley MT, Mooney KM. Computational systems biology for aging research, *Interdiscip Top Gerontol* 2015;40:35-48.
48. Mc Auley MT, Mooney KM. Computationally Modeling Lipid Metabolism and Aging: A Mini-review, *Computational and Structural Biotechnology Journal* 2015;13:38-46.
49. Mc Auley MT, Mooney KM, Angell PJ et al. Mathematical Modelling of Metabolic Regulation in Aging, *Metabolites* 2015, 5(2), 232-251; 2015;5:232-251.
50. Mc Auley MT, Proctor CJ, Corfe BM et al. Nutrition Research and the Impact of Computational Systems Biology, *Journal of Computer Science and Systems Biology* 2013;6:271-285.
51. Robertson KD. DNA methylation, methyltransferases, and cancer, *Oncogene* 2001;20:3139-3155.
52. Arasaradnam RP, Commane DM, Bradburn D et al. A review of dietary factors and its influence on DNA methylation in colorectal carcinogenesis, *Epigenetics* 2008;3:193-198.
53. Bailey LB, Gregory JF, 3rd. Folate metabolism and requirements, *J Nutr* 1999;129:779-782.
54. Robertson KD, Uzvolgyi E, Liang G et al. The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors, *Nucleic Acids Res* 1999;27:2291-2298.
55. Chen T, Ueda Y, Dodge JE et al. Establishment and maintenance of genomic methylation patterns in mouse embryonic stem cells by Dnmt3a and Dnmt3b, *Mol Cell Biol* 2003;23:5594-5605.
56. Okano M, Bell DW, Haber DA et al. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development, *Cell* 1999;99:247-257.
57. Scourzic L, Mouly E, Bernard OA. TET proteins and the control of cytosine demethylation in cancer, *Genome Med* 2015;7:9.
58. Landan G, Cohen NM, Mukamel Z et al. Epigenetic polymorphism and the stochastic formation of differentially methylated regions in normal and cancerous tissues, *Nat Genet* 2012;44:1207-1214.
59. Shipony Z, Mukamel Z, Cohen NM et al. Dynamic and static maintenance of epigenetic memory in pluripotent and somatic cells, *Nature* 2014;513:115-119.
60. Nijhout HF, Reed MC, Ulrich CM. Mathematical models of folate-mediated one-carbon metabolism, *Vitam Horm* 2008;79:45-82.
61. Voit EO, Martens HA, Omholt SW. 150 years of the mass action law, *PLoS Comput Biol* 2015;11:e1004012.
62. Michaelis L, Menten ML, Johnson KA et al. The original Michaelis constant: translation of the 1913 Michaelis-Menten paper, *Biochemistry* 2011;50:8264-8269.


63. Enrique Salcedo-Sora J, Mc Auley MT. A mathematical model of microbial folate biosynthesis and utilisation: implications for antifolate development, *Mol Biosyst* 2016;12:923-933.
64. Jackson RC, Harrap KR. Studies with a mathematical model of folate metabolism, *Arch Biochem Biophys* 1973;158:827-841.
65. Nijhout HF, Reed MC, Budu P et al. A mathematical model of the folate cycle: new insights into folate homeostasis, *J Biol Chem* 2004;279:55008-55016.
66. Reed MC, Nijhout HF, Neuhauser ML et al. A mathematical model gives insights into nutritional and genetic aspects of folate-mediated one-carbon metabolism, *J Nutr* 2006;136:2653-2661.
67. Reed MC, Nijhout HF, Sparks R et al. A mathematical model of the methionine cycle, *J Theor Biol* 2004;226:33-43.
68. Nijhout HF, Reed MC, Anderson DF et al. Long-range allosteric interactions between the folate and methionine cycles stabilize DNA methylation reaction rate, *Epigenetics* 2006;1:81-87.
69. Neuhauser ML, Nijhout HF, Gregory JF, 3rd et al. Mathematical modeling predicts the effect of folate deficiency and excess on cancer-related biomarkers, *Cancer Epidemiol Biomarkers Prev* 2011;20:1912-1917.
70. Duncan TM, Reed MC, Nijhout HF. A population model of folate-mediated one-carbon metabolism, *Nutrients* 2013;5:2457-2474.
71. Capra JA, Kostka D. Modeling DNA methylation dynamics with approaches from phylogenetics, *Bioinformatics* 2014;30:i408-414.
72. Genreux DP, Miner BE, Bergstrom CT et al. A population-epigenetic model to infer site-specific methylation rates from double-stranded DNA methylation patterns, *Proc Natl Acad Sci U S A* 2005;102:5802-5807.
73. Flottmann M, Scharp T, Klipp E. A stochastic model of epigenetic dynamics in somatic cell reprogramming, *Front Physiol* 2012;3:216.
74. Zheng H, Wu H, Li J et al. CpGIMethPred: computational model for predicting methylation status of CpG islands in human genome, *BMC Med Genomics* 2013;6 Suppl 1:S13.
75. Feinberg AP, Irizarry RA. Evolution in health and medicine Sackler colloquium: Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease, *Proc Natl Acad Sci U S A* 2010;107 Suppl 1:1757-1764.
76. Przybilla J, Rohlf T, Loeffler M et al. Understanding epigenetic changes in aging stem cells--a computational model approach, *Aging Cell* 2014;13:320-328.
77. Ro S, Rannala B. Methylation patterns and mathematical models reveal dynamics of stem cell turnover in the human colon, *Proc Natl Acad Sci U S A* 2001;98:10519-10521.
78. Otto SP, Walbot V. DNA methylation in eukaryotes: kinetics of demethylation and de novo methylation during the life cycle, *Genetics* 1990;124:429-437.
79. Sontag LB, Lorincz MC, Georg Luebeck E. Dynamics, stability and inheritance of somatic DNA methylation imprints, *J Theor Biol* 2006;242:890-899.
80. McGovern AP, Powell BE, Chevassut TJ. A dynamic multi-compartmental model of DNA methylation with demonstrable predictive value in hematological malignancies, *J Theor Biol* 2012;310:14-20.
81. Pfeifer GP, Steigerwald SD, Hansen RS et al. Polymerase chain reaction-aided genomic sequencing of an X chromosome-linked CpG island: methylation patterns suggest clonal inheritance, CpG site autonomy, and an explanation of activity state stability, *Proc Natl Acad Sci U S A* 1990;87:8252-8256.
82. Riggs AD, Xiong Z. Methylation and epigenetic fidelity, *Proc Natl Acad Sci U S A* 2004;101:4-5.

83. Jeltsch A, Jurkowska RZ. New concepts in DNA methylation, *Trends Biochem Sci* 2014;39:310-318.
84. Haerter JO, Lovkvist C, Dodd IB et al. Collaboration between CpG sites is needed for stable somatic inheritance of DNA methylation states, *Nucleic Acids Res* 2014;42:2235-2244.
85. Olariu V, Lovkvist C, Sneppen K. Nanog, Oct4 and Tet1 interplay in establishing pluripotency, *Sci Rep* 2016;6:25438.
86. Pufulete M, Al-Ghnaniem R, Leather AJ et al. Folate status, genomic DNA hypomethylation, and risk of colorectal adenoma and cancer: a case control study, *Gastroenterology* 2003;124:1240-1248.
87. Weiner AS, Boyarskikh UA, Voronina EN et al. Methylenetetrahydrofolate reductase C677T and methionine synthase A2756G polymorphisms influence on leukocyte genomic DNA methylation level, *Gene* 2014;533:168-172.
88. Harmon DL, Shields DC, Woodside JV et al. Methionine synthase D919G polymorphism is a significant but modest determinant of circulating homocysteine concentrations, *Genet Epidemiol* 1999;17:298-309.
89. Ganguly P, Alam SF. Role of homocysteine in the development of cardiovascular disease, *Nutr J* 2015;14:6.
90. Zhang JG, Liu JX, Li ZH et al. Dysfunction of endothelial NO system originated from homocysteine-induced aberrant methylation pattern in promoter region of DDAH2 gene, *Chin Med J (Engl)* 2007;120:2132-2137.
91. Bonsch D, Lenz B, Reulbach U et al. Homocysteine associated genomic DNA hypermethylation in patients with chronic alcoholism, *J Neural Transm (Vienna)* 2004;111:1611-1616.
92. Zhi Y, Huang Y, Li Z et al. [Hypermethylation of estrogen receptor-alpha gene and high homocysteine in atheromatosis patients], *Wei Sheng Yan Jiu* 2008;37:314-317.
93. Huang Y, Peng K, Su J et al. Different effects of homocysteine and oxidized low density lipoprotein on methylation status in the promoter region of the estrogen receptor alpha gene, *Acta Biochim Biophys Sin (Shanghai)* 2007;39:19-26.
94. Lee ME, Wang H. Homocysteine and hypomethylation. A novel link to vascular disease, *Trends Cardiovasc Med* 1999;9:49-54.
95. Yi P, Melnyk S, Pogribna M et al. Increase in plasma homocysteine associated with parallel increases in plasma S-adenosylhomocysteine and lymphocyte DNA hypomethylation, *J Biol Chem* 2000;275:29318-29323.
96. Zhang D, Sun X, Liu J et al. Homocysteine accelerates senescence of endothelial cells via DNA hypomethylation of human telomerase reverse transcriptase, *Arterioscler Thromb Vasc Biol* 2015;35:71-78.
97. MacFarlane AJ, Liu X, Perry CA et al. Cytoplasmic serine hydroxymethyltransferase regulates the metabolic partitioning of methylenetetrahydrofolate but is not essential in mice, *J Biol Chem* 2008;283:25846-25853.
98. Herbig K, Chiang EP, Lee LR et al. Cytoplasmic serine hydroxymethyltransferase mediates competition between folate-dependent deoxyribonucleotide and S-adenosylmethionine biosyntheses, *J Biol Chem* 2002;277:38381-38389.
99. Scotti M, Stella L, Shearer EJ et al. Modeling cellular compartmentation in one-carbon metabolism, *Wiley Interdiscip Rev Syst Biol Med* 2013;5:343-365.
100. Field MS, Kamynina E, Agunloye OC et al. Nuclear enrichment of folate cofactors and methylenetetrahydrofolate dehydrogenase 1 (MTHFD1) protect de novo thymidylate biosynthesis during folate deficiency, *J Biol Chem* 2014;289:29642-29650.


101. Groth M, Moissiard G, Wirtz M. MTHFD1 controls DNA methylation in *Arabidopsis* 2016;7:11640.
102. Anderson DD, Woeller CF, Chiang EP et al. Serine hydroxymethyltransferase anchors de novo thymidylate synthesis pathway to nuclear lamina for DNA synthesis, *J Biol Chem* 2012;287:7051-7062.
103. Anderson DD, Eom JY, Stover PJ. Competition between sumoylation and ubiquitination of serine hydroxymethyltransferase 1 determines its nuclear localization and its accumulation in the nucleus, *J Biol Chem* 2012;287:4790-4799.
104. Lee B, Muller MT. SUMOylation enhances DNA methyltransferase 1 activity, *Biochem J* 2009;421:449-461.
105. Ling Y, Sankpal UT, Robertson AK et al. Modification of de novo DNA methyltransferase 3a (Dnmt3a) by SUMO-1 modulates its interaction with histone deacetylases (HDACs) and its capacity to repress transcription, *Nucleic Acids Res* 2004;32:598-610.
106. Gutenkunst RN, Waterfall JJ, Casey FP et al. Universally sloppy parameter sensitivities in systems biology models, *PLoS Comput Biol* 2007;3:1871-1878.
107. Scheer M, Grote A, Chang A et al. BRENDA, the enzyme information system in 2011, *Nucleic Acids Res* 2011;39:D670-676.
108. Wittig U, Kania R, Golebiewski M et al. SABIO-RK--database for biochemical reaction kinetics, *Nucleic Acids Res* 2012;40:D790-796.
109. Pradhan M, Esteve PO, Chin HG et al. CXXC domain of human DNMT1 is essential for enzymatic activity, *Biochemistry* 2008;47:10000-10009.
110. Balada E, Ordi-Ros J, Serrano-Acedo S et al. Transcript levels of DNA methyltransferases DNMT1, DNMT3A and DNMT3B in CD4+ T cells from patients with systemic lupus erythematosus, *Immunology* 2008;124:339-347.
111. Gibson MA, Bruck J. Efficient exact stochastic simulation of chemical systems with many species and many channels. , *J. Phys. Chem* 2000;A 104:1876-1889.
112. Gillespie D. A general method for numerically simulating stochastic time evolution of coupled chemical-reactions., *Journal of Computational Physics* 1976;22::403-434.
113. Elowitz MB, Levine AJ, Siggia ED et al. Stochastic gene expression in a single cell, *Science* 2002;297:1183-1186.
114. Singh A, Soltani M. Quantifying intrinsic and extrinsic variability in stochastic gene expression models, *PLoS One* 2013;8:e84301.
115. Raser JM, O'Shea EK. Control of stochasticity in eukaryotic gene expression, *Science* 2004;304:1811-1814.
116. Raj A, van Oudenaarden A. Nature, nurture, or chance: stochastic gene expression and its consequences, *Cell* 2008;135:216-226.
117. Wilkinson DJ. Stochastic modelling for quantitative description of heterogeneous biological systems, *Nat Rev Genet* 2009;10:122-133.
118. Wei T, Dai H. Quantification of GFP signals by fluorescent microscopy and flow cytometry, *Methods Mol Biol* 2014;1163:23-31.
119. Soboleski MR, Oaks J, Halford WP. Green fluorescent protein is a quantitative reporter of gene expression in individual eukaryotic cells, *Faseb j* 2005;19:440-442.
120. Sigal A, Milo R, Cohen A et al. Variability and memory of protein levels in human cells, *Nature* 2006;444:643-646.
121. Thattai M, van Oudenaarden A. Intrinsic noise in gene regulatory networks, *Proc Natl Acad Sci U S A* 2001;98:8614-8619.
122. Hansen KD, Timp W, Bravo HC et al. Increased methylation variation in epigenetic domains across cancer types, *Nat Genet* 2011;43:768-775.

123. Cheow LF, Courtois ET, Tan Y et al. Single-cell multimodal profiling reveals cellular epigenetic heterogeneity, *Nat Methods* 2016.
124. Proctor CJ, Gray DA. Explaining oscillations and variability in the p53-Mdm2 system, *BMC Syst Biol* 2008;2:75.
125. Wu J, Vidakovic B, Voit EO. Constructing stochastic models from deterministic process equations by propensity adjustment, *BMC Syst Biol* 2011;5:187.
126. Ball DA, Ahn TH, Wang P et al. Stochastic exit from mitosis in budding yeast: model predictions and experimental observations, *Cell Cycle* 2011;10:999-1009.
127. Puchalka J, Kierzek AM. Bridging the gap between stochastic and deterministic regimes in the kinetic simulations of the biochemical reaction networks, *Biophys J* 2004;86:1357-1372.
128. McEntee G, Minguzzi S, O'Brien K et al. The former annotated human pseudogene dihydrofolate reductase-like 1 (DHFR1L1) is expressed and functional, *Proc Natl Acad Sci U S A* 2011;108:15157-15162.
129. Christensen KE, Mirza IA, Berghuis AM et al. Magnesium and phosphate ions enable NAD binding to methylenetetrahydrofolate dehydrogenase-methenyltetrahydrofolate cyclohydrolase, *J Biol Chem* 2005;280:34316-34323.
130. Burda P, Schafer A, Suormala T et al. Insights into severe 5,10-methylenetetrahydrofolate reductase deficiency: molecular genetic and enzymatic characterization of 76 patients, *Hum Mutat* 2015;36:611-621.
131. Smith GK, Banks SD, Monaco TJ et al. Activity of an NAD-dependent 5,10-methylenetetrahydrofolate dehydrogenase in normal tissue, neoplastic cells, and oncogene-transformed cells, *Arch Biochem Biophys* 1990;283:367-371.
132. Wolthers KR, Scrutton NS. Cobalamin uptake and reactivation occurs through specific protein interactions in the methionine synthase-methionine synthase reductase complex, *FEBS J* 2009;276:1942-1951.
133. Kotb M, Kredich NM. S-Adenosylmethionine synthetase from human lymphocytes. Purification and characterization, *J Biol Chem* 1985;260:3923-3930.
134. Bacolla A, Pradhan S, Larson JE et al. Recombinant human DNA (cytosine-5) methyltransferase. III. Allosteric control, reaction order, and influence of plasmid topology and triplet repeat length on methylation of the fragile X CGG.CCG sequence, *J Biol Chem* 2001;276:18605-18613.
135. Pfeifer GP, Grunwald S, Boehm TL et al. Isolation and characterization of DNA cytosine 5-methyltransferase from human placenta, *Biochim Biophys Acta* 1983;740:323-330.
136. Roy PH, Weissbach A. DNA methylase from HeLa cell nuclei, *Nucleic Acids Res* 1975;2:1669-1684.
137. Bosviel R, Garcia S, Lavediaux G et al. BRCA1 promoter methylation in peripheral blood DNA was identified in sporadic breast cancer and controls, *Cancer Epidemiol* 2012;36:e177-182.
138. Hunter A, Spechler PA, Cwanger A et al. DNA methylation is associated with altered gene expression in AMD, *Invest Ophthalmol Vis Sci* 2012;53:2089-2105.
139. Kwabi-Addo B, Chung W, Shen L et al. Age-related DNA methylation changes in normal human prostate tissues, *Clin Cancer Res* 2007;13:3796-3802.
140. Nakagawa H, Nuovo GJ, Zervos EE et al. Age-related hypermethylation of the 5' region of MLH1 in normal colonic mucosa is associated with microsatellite-unstable colorectal cancer development, *Cancer Res* 2001;61:6991-6995.
141. Kawakami K, Ruzskiewicz A, Bennett G et al. DNA hypermethylation in the normal colonic mucosa of patients with colorectal cancer, *Br J Cancer* 2006;94:593-598.


142. Scott JL, Gabrielides C, Davidson RK et al. Superoxide dismutase downregulation in osteoarthritis progression and end-stage disease, *Ann Rheum Dis* 2010;69:1502-1510.
143. Polosak J, Kurylowicz A, Roszkowska-Gancarz M et al. Aging is accompanied by a progressive decrease of expression of the WRN gene in human blood mononuclear cells, *J Gerontol A Biol Sci Med Sci* 2011;66:19-25.
144. Takami N, Osawa K, Miura Y et al. Hypermethylated promoter region of DR3, the death receptor 3 gene, in rheumatoid arthritis synovial cells, *Arthritis Rheum* 2006;54:779-787.
145. Loeser RF, Im HJ, Richardson B et al. Methylation of the OP-1 promoter: potential role in the age-related decline in OP-1 expression in cartilage, *Osteoarthritis Cartilage* 2009;17:513-517.




(a) Defining promoter/gene body methylation using intrinsic & extrinsic noise



Young Partially Methylated CpG Islands



Age



Old Hypermethylation

(7) Promoter methylation can accompany gene silencing

(b) Indicative reaction list for a combined system which could be modelled as propensity functions

F $\xrightarrow{1}$ DHF
 DHF $\xrightarrow{1}$ THF
 THF $\xrightarrow{1}$ 510MTHF
 510MTHF $\xrightarrow{1}$ 5MTHF
 510MTHF $\xrightarrow{1}$ DHF
 5MTHF+Hcy $\xrightarrow{1}$ THF +Met
 Met $\xrightarrow{1}$ SAM
 THF $\xrightarrow{1}$ 10FORM
 10FORM $\xrightarrow{1}$ DHF
 SAM $\xrightarrow{1}$ CpGCH₃+SAH
 SAH $\xrightarrow{1}$ Hcy
 10FORM+510MTHF $\xrightarrow{1}$ DNA
 CpGCH₃ $\xrightarrow{1}$ CpG
 CpGCH₃ $\xrightarrow{1}$ CpG

Genes
BRAC1
IGF2
GSTM1
GSTP1
MLH1
SOD2
WRN
ER
DR3

Decrease d \rightarrow expression

Table 1 A summary of the models of folate metabolism cited in the text.

Authors	Title	Model Overview/Predictions	Computational Methods Used
Jackson and Harrap (1973) [64]	Studies with a mathematical model of folate metabolism	Included the main reactions involved in cellular folate metabolism in mouse leukaemia cells. Used to predict time course changes in purine and thymidylate biosynthesis rates.	A deterministic framework consisting of a system of integrated equations. Kinetic parameters were determined from pre-existing experimental data.
Nijhout et al. (2004) [65]	A mathematical model of the folate cycle: new insights into folate homeostasis	Folate cycle underpinned by enzyme kinetics. Used to explore the 'methyl trap hypothesis', the effects of methotrexate and sensitivities to variation in enzymatic parameters	A deterministic ODE based kinetic model constructed using the assumption that most reactions in the folate cycle are bimolecular.
Reed et al. (2006) [66]	A mathematical model gives insights into nutritional and genetic aspects of folate-mediated one-carbon metabolism	Model of FOCM. Predictions included the inverse association between hcy and very low folate levels, DNA methylation reaction rate is relatively insensitive to changes in folate pool size, folate concentrations become very high, enzyme velocities diminish, mutations in MTHFR reduces SAM	Deterministic ODE model created by combining the model by Nijhout et al. (2004) and models of methionine metabolism [67, 68]
Neuhouser et al. (2011) [69]	Mathematical modeling predicts the effect of folate deficiency and excess on cancer-related biomarkers	FOCM model used by this group. Used to predict: the effect of a broad range of intracellular folate concentrations simulating variation in folate status on mechanisms relevant to carcinogenesis.	Deterministic model based on [66].
Duncan et al. (2013) [70]	A Population Model of Folate-Mediated One-Carbon Metabolism[70]	Compartmental body model folate metabolism. Showed tissue and plasma folate is highly correlated, but liver and plasma folate much less so. Oxidative stress increases the plasma SAM/SAH ratio.	Models from [65, 66] were incorporated into liver and plasma compartment. Coupled set of ODEs Random number used to generate a virtual population of 10000 individual each with different folate metabolism Model constructed using Matlab
Salcedo-Sora and Mc Auley (2016) [63]	A mathematical model of microbial folate biosynthesis and utilisation: implications for antifolate development	First model to be assembled of folate biosynthesis and utilisation. The model suggested a particular small set of folate intermediates are specific targets which could complement current antifolates, and the model substantiates the case for a substrate cycle in the folinic acid biosynthesis reaction.	Deterministic ODE model Folate pathway informed by the KEGG database http://www.genome.jp/kegg/) and kinetic parameters were compile from the enzyme database BRENDA http://www.brenda-enzymes.org/

Table 2 A summary of DNA methylation models cited in the text.

Authors	Title	Model Features /Predictions	Computational Methods Used
Capra <i>et al.</i> (2014) [71]	Modeling DNA methylation dynamics with approaches from phylogenetics	The model was capable of inferring unobserved CpG methylation states from observations at the same sites in related cell types (90% correct).	Continuous-time Markov chain approach. R source code: www.kostkalab.net/software .
Genereux <i>et al.</i> (2005) [72]	A population-epigenetic model to infer site-specific methylation rates from double-stranded DNA methylation patterns.	Population-model of DNA methylation dynamics. Can infer site-specific rates of both maintenance and <i>de novo</i> methylation values.	Probabilistic model designed to track the frequencies of methylated, hemimethylated, and unmethylated CpG/CpG dyads at a given CpG site across a population of cells from a single tissue of a single individual.
Flottmann <i>et al.</i> (2012) [73]	A stochastic model of epigenetic dynamics in somatic cell reprogramming	Describes the interplay between gene expression, chromatin modifications, and DNA methylation. Infers cytosine methylation rates at several sites within the promoter of the human gene <i>FMR1</i> .	Probabilistic Boolean networks to represent two kinds of stochasticity.
Zheng <i>et al.</i> (2013) [74]	CpGIMethPred : computational model for predicting methylation status of CpG islands in human genome	Machine-based models used to predict the methylation status of the CpG islands in the human genome under normal conditions.	Three part model systems to obtain CpG island map. This involved database extraction of methylation data, feature extraction and statistical predictive modelling.
Feinberg <i>et al.</i> (2010) [75]	Evolution in health and medicine Sackler colloquium: Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease	Epigenetic evolutionary model. Provided evidence for stochastic epigenetic variation, a heritable genetic mechanism for variable methylation, and genetically inherited stochastic variation in evolution, in changing environments mediated epigenetically.	Expanded the Fisher-Wright neutral selection model.
Przybilla <i>et al.</i> (2014) [76]	Understanding epigenetic changes in aging stem cells--a computational model	Used a model to explore age-related changes in DNA methylation within stem cells and simulations suggested homing at stem cell niches retarded epigenetic ageing.	Utilised the model of DNA by Sontag <i>et al.</i> (2006) [79]

	approach		
Yatabe <i>et al.</i> (2001) [77]	Methylation patterns and mathematical models reveal dynamics of stem cell turnover in the human colon	Experimental work and simulations suggest human crypts are long-lived, accumulate random methylation errors, and contain multiple stem cells that go through “bottlenecks” during life.	Stochastic model. Crypts were simulated with a methylation error rate of 2×10^{-5} per CpG site per division, and one division per day.
Otto <i>et al.</i> (1990) [78]	DNA methylation in eukaryotes: kinetics of demethylation and de novo methylation during the life cycle.	Steady state/equilibrium achieved such that the proportion of sites which are newly methylated equals the proportion of sites which become demethylated in a cell generation.	Kinetic model that represents methylation and demethylation. It incorporates values for <i>de novo</i> methylation and the error rate equation based model that ignores randomness.
Sontag <i>et al.</i> (2006) [79]	Dynamics, stability and inheritance of somatic DNA methylation imprints.	Describes the evolution of hypo- to hypermethylated equilibria as a result of methylation noise in a finite system of CpG sites.	Markov-chain model, CpGs can change methylation states during cell division only based on probabilities for conservation/maintenance/loss and <i>de novo</i> methylation.
McGovern <i>et al.</i> (2012) [80]	A dynamic multi-compartmental model of DNA methylation with demonstrable predictive value in hematological malignancies.	Was able to fully represent the full suite of DNA methylation/demethylation reactions and was used as a tool for predicting haematological malignancies	Constructed with partial differential equations.
Riggs and Xiong (2004) [82]	Methylation and epigenetic fidelity	Methylation at most sites in cell lines, and in tissue cell types is stochastically variable to a certain extent, with each site depending on site-specific probabilities.	The stochastic methylation model assumes that for each CpG dyad in each DNA molecule there is a certain efficiency (probability) of methylation maintenance failure or <i>de novo</i> methylation.
Jeltsch and Jurkowska (2014) [83]	New concepts in DNA methylation	Added to the Riggs model [82] by including the rate of cell division as well as active and passive demethylation.	Stochastic equation model with rates of cell division and rates of active and passive demethylation.
Haerter <i>et al.</i> (2014) [84]	Collaboration between CpG sites is needed for stable somatic inheritance of DNA methylation	Compared the standard/classical model to a collaborative model and showed that the standard model is inconsistent with many experimental observations.	Stochastic model implemented using the Gillespie algorithm [112].

	states		
Olariu et al. (2016) [85]	Nanog, Oct4 and Tet1 interplay in establishing pluripotency.	Assembled a regulatory network model of Oct4, Nanog and Tet1 which included positive feedback loops involving DNA-demethylation around the promoters of Oct4 and Tet1. Provides a template for novel framework combining transcription regulation with DNA methylation modifications	To model methylation the stochastic model by Haerter et al. (2014) [84] was used.

Table 3. Examples of kinetic parameters available for assembly of a deterministic model

Authors	Reaction	Parameters identified	Cell line	Monoglutamated or polyglutamated	Study Details
McEntee <i>et al.</i> (2011) [128].	Dihydrofolate-> Tetrahydrofolate	$V_{max} = 0.0132 \mu\text{m}/\text{min}/\text{mL}$ $K_m = 20.1 \mu\text{M}$	Human cell line cell line BT474.	Polyglutamated *	The kinetics of the dihydrofolate reductase enzyme were compared to a similar enzyme in order to determine its parameters.
Christensen <i>et al.</i> (2005) [129].	Tetrahydrofolate-> 5,10-Methylenetetrahydrofolate	$V_{max} = 22.5 \mu\text{m}/\text{min}/\text{mL}$ $K_m = 412 \mu\text{M}$	Human homology model.	Polyglutamated	The kinetics of this reaction were determined using a human homology model of NAD-dependent methylenetetrahydrofolate dehydrogenase-cyclohydrolase.
Burda <i>et al.</i> (2015) [130] Smith <i>et al.</i> (1990) [131].	5,10-Methylenetetrahydrofolate->5-methyltetrahydrofolate	$K_m = 29.95 \mu\text{M}$ Specific activity = $0.005 \mu\text{mol}/\text{min}/\text{mg}$	Human skin fibroblasts [130] and human liver cells for specific activity [131]	Polyglutamated	Mean k_m value taken from [130]. Specific activity taken from analysis of human liver [131].
Wolthers and Scrutton (2009) [132].	5-methyltetrahydrofolate+ Homocysteine ->Methionine +Tetrahydrofolate	$K_m = 29.95 \mu\text{M}$ Specific activity = $1.54 \mu\text{mol}/\text{min}/\text{mg}$	Human MS derived from <i>Pichia pastoris</i> .	Polyglutamated	The kinetics of human methionine synthase were identified by its expression and purification from <i>Pichia pastoris</i> .
Kotb and Kredich (1985) [133].	Methionine-> s-adenosyl methionine	$K_m = 3.3 \mu\text{M}$ Specific activity = $12.2 \mu\text{mol}/\text{min}/\text{mg}$	Human lymphocytes	Polyglutamated	Adenosylmethionine synthetase was purified from human lymphocytes.
Bacolla <i>et al.</i> (2001) [134] Pfeife <i>et al.</i> (1983) [135].	DNA+s-adenosyl methionine-> s-adenosyl homocysteine + CpGCH ₃ (maintenance reaction)	$K_m^{\text{Aldomet}} = 11 \mu\text{M}$ $K_m^{\text{DNA}} = 23 \mu\text{M}$ Specific activity = $0.000016 \mu\text{mol}/\text{min}/\text{mg}$	Human constructs[134] And human placenta [135].	Polyglutamated	Steady-state kinetic analyses of human DNMT1 [134] Isolation and characterization of DNA cytosine 5-methyltransferase from human placenta [135].
Roy and Weissbach (1975) [136].	CpGCH ₃ -> CpG (<i>de novo</i> reaction)	$K_m^{\text{Aldomet}} = 0.00325 \mu\text{M}$ $K_m^{\text{DNA}} = 0.086 \mu\text{M}$ Specific activity = $0.000013 \mu\text{mol}/\text{min}/\text{mg}$	HeLa cells	Polyglutamated	Enzymes were purified from HeLa cell nuclei by chromatography on diethylaminoethyl cellulose.
	CpG-> CpGCH ₃ (Active Demethylation)	Unidentified			

*Although the studies do not state explicitly the chemical form of the folates, as they were cell line assays it can be assumed the chemical administered to the cells was monoglutamated. However, it is likely to have become polyglutamated by the time the enzymatic assay took place.

Table 4. Examples of Genes associated ageing/health-span which have been found to be hypermethylated with age

Gene	Full name	Function	Study and Conclusions*	Source
BRAC1	Breast cancer 1	DNA repair	Cancer study: Both controls and cancer patients displayed methylation of BRAC1	Bosviel <i>et al.</i> (2012) [137]
IGF2	Insulin like growth factor II	Promotes cell growth and division.	Cancer study: methylation change in IGF2 promoter during aging and carcinogenesis.	Issa <i>et al.</i> (1996) [24]
GSTM1	Glutathione S-transferase isoform mu1 (GSTM1)	Codes for an enzyme that has a role to play in the prevention of oxidative damage	GSTM1 promoter methylation was confirmed by CpG island methylation in age related macular degeneration subjects.	Hunter <i>et al.</i> (2012) [138]
GSTP1	Glutathione S-transferase P	Codes for an enzyme that has an important role to play in the prevention of oxidative damage.	Cancer study: Changes in Normal Human Prostate Tissues. Increase in promoter methylation with age for CpG islands in normal prostate tissue samples in several genes.	Kwabi-Addo <i>et al.</i> (2007) [139]
MLH1	MutL homolog 1	Codes for a protein involved in DNA mismatch repair	Cancer study: found partially methylated alleles in 44% of patients <60yrs and in 83% ≥ 80 yrs. Fully methylated alleles in 55% of patients with Microsatellite instability + tumours.	Nakagawa <i>et al.</i> (2001) [140]
			Cancer study: found hypermethylation in the normal colonic mucosa of patients with colorectal cancer. Methylation of CpG islands is seen in the normal colonic mucosa and increased with age.	Kawakami <i>et al.</i> (2006) [141]
SOD2		Codes for an enzyme that helps prevent oxidative damage	Osteoarthritis (OA) study: used human and guinea pig tissue samples. SOD2 promoter had significant DNA methylation alterations in OA cartilage.	Scott <i>et al.</i> (2010) [142]
WRN	Werner syndrome	Codes for a DNA helicase that participates in genome maintenance	Immunosenescence study: Analysis of the 361 bp WRN promoter CpG island showed a slight increase in methylation with age of its methylation status. Authors report that it was biologically negligible.	Polosak <i>et al.</i> (2011) [143]
ER	Estrogen receptor	Estrogen receptor	Cancer study: CpG island methylation, of ER gene in a subpopulation of cells increased as a direct function of age in human colonic mucosa.	Issa <i>et al.</i> (1994)[27]

DR3	The death receptor 3 (DR3) gene	Member of the apoptosis-inducing fas receptor gene family	Rheumatoid Arthritis (RA) study: Hypermethylated promoter region of DR3, the death receptor 3 gene, in rheumatoid arthritis synovial cells.	Takami <i>et al.</i> (2006) [144]
(OP-1, BMP-7)	Oestrogenic protein 1	Bone maintenance	Osteoporosis study: Methylation of the OP-1 promoter was found in chondrocytes from tissue obtained from older adults (study age range 19-86yrs).	Loeser <i>et al.</i> (2009) [145]