

Minkowski medial axes and shocks of plane curves

Graham Mark Reeve* and Farid Tari†

Abstract

In this paper a Minkowski analogue of the Euclidean medial axis of a closed and smooth plane curve is introduced. We study its generic local configurations and determine the types of shocks that can occur on these.

1 Introduction

The concept of the *medial axis* of Euclidean plane curves was first introduced by Blum in [1]. For a closed and smooth curve γ the *medial axis* is defined to be the locus of the centres of maximal circles that are tangent to γ in two or more points. Here a circle is said to be *maximal* if its radius equals the absolute minimum distance from its centre to γ : such a circle is either contained in the interior or the exterior of γ and cannot be expanded about its centre without crossing γ . Many applications of medial axes are given in Blum's original paper [1] and other applications relating to computer vision can be found in [10].

The *symmetry set* of the curve γ is the same as that of the medial axis except that the constraint that the circles must be maximal is dropped (see [4, 5]). For this reason the medial axis forms a subset of the symmetry set.

In [6] the generic shocks that can occur on the Euclidean medial axis are classified. Motivated by applications in fluid mechanics, Bogaevsky uses a different approach in [3] to obtain similar results to those in [6].

*Supported by the FAPESP grant 2012/0536-9.

†Partially supported by the grants FAPESP 2011/21240-4, CNPq 301589/2012-7 and CNPq 472796/2013-5.

2000 Mathematics Subject classification. Primary 53A35 ; Secondary 58K05 53D12

Key Words and Phrases. Curves, Minkowski plane, Medial axis, Shocks, Singularities, Minkowski symmetry sets.

In this paper we introduce a Minkowski analogue of the Euclidean medial axis for a closed and smooth plane curve, called the Minkowski medial axis (*MMA*). An analogue of the symmetry set for curves in the Minkowski plane, called the *Minkowski Symmetry Set* (*MSS*), is introduced and studied in [11]. For a curve γ in the Minkowski plane, the *MSS* is defined as the locus of the centres of pseudo-circles that are tangent to the curve γ in two or more points.

Similarly to the Euclidean version, a point on the *MSS* is said to belong to the Minkowski medial axis if the radius r of the bi-tangent pseudo-circle equals the absolute maximum (if r is positive) or the absolute minimum (if r is negative) distance from its centre to γ . We show in Theorem 4.7 that this definition implies that the bi-tangent points lie on just one branch of the pseudo-circle. This fact leads on to a new generalised type of *MMA*, called the *1-branch MMA*. This is defined to be the centres of pseudo-circles that have a bitangency with just one branch of the pseudo-circles. It follows that the *MMA* forms a subset of the *1-branch MMA*. The Minkowski symmetry set together with a radius function, like its Euclidean counterpart ([10]), can be used to reconstruct the original curve γ . However, unlike with the Euclidean medial axis, neither the *MMA* nor the *1-branch MMA* can be used to reconstruct *non-convex* curves. However, the *1-branch MMA* together with a radius function can be used to reconstruct *convex* curves.

The *shock set* of γ is obtained by adding an arrow to the *MMA* indicating the direction of increasing radii of the relevant bi-tangent pseudo-circles. In this paper, we obtain the generic local configuration of the *MMA* and shocks of curves in the Minkowski plane.

2 Preliminaries

The *Minkowski plane* $(\mathbb{R}_1^2, \langle, \rangle)$ is the vector space \mathbb{R}^2 endowed with the pseudo-scalar product $\langle \mathbf{u}, \mathbf{v} \rangle = -u_0v_0 + u_1v_1$, for any $\mathbf{u} = (u_0, u_1)$ and $\mathbf{v} = (v_0, v_1)$. A vector $\mathbf{u} \in \mathbb{R}_1^2 \setminus \{0\}$ is called

- spacelike* if $\langle \mathbf{u}, \mathbf{u} \rangle > 0$,
- timelike* if $\langle \mathbf{u}, \mathbf{u} \rangle < 0$ or
- lightlike* if $\langle \mathbf{u}, \mathbf{u} \rangle = 0$.

The norm of a vector is defined by $\|\mathbf{u}\| = \sqrt{|\langle \mathbf{u}, \mathbf{u} \rangle|}$.

The pseudo-circles in \mathbb{R}_1^2 with centre $c \in \mathbb{R}_1^2$ and radius r are defined as follows:

$$\begin{aligned} H^1(c, r) &= \{p \in \mathbb{R}_1^2 \mid \langle p - c, p - c \rangle = -r^2\} \text{ if } r < 0, \\ S_1^1(c, r) &= \{p \in \mathbb{R}_1^2 \mid \langle p - c, p - c \rangle = r^2\} \text{ if } r > 0, \\ LC^*(c) &= \{p \in \mathbb{R}_1^2 \setminus 0 \mid \langle p - c, p - c \rangle = 0\} \text{ if } r = 0. \end{aligned}$$

Observe that $LC^*(c)$ is the union of the two lines through c with tangent directions $(1, 1)$ and $(1, -1)$, with the point c removed. The pseudo-circle $H^1(c, -r)$




Figure 1: The three types of vectors (left) and pseudo-circles (right) in \mathbb{R}_1^2 .

has two branches which can be parametrised by $c + (\pm r \cosh(t), r \sinh(t))$, $t \in \mathbb{R}$. The pseudo-circle $S^1(c, r)$ also has two branches which can be parametrised by $c + (r \sinh(t), \pm r \cosh(t))$, $t \in \mathbb{R}$. See Figure 1.

Let $\gamma : J \rightarrow \mathbb{R}_1^2$ be a smooth curve, where J is an open interval of \mathbb{R} or the unit circle S^1 if the curve is closed. The curve γ is spacelike if $\gamma'(t)$ is a spacelike vector for all $t \in J$ and is timelike if $\gamma'(t)$ is a timelike vector for all $t \in J$. A point $\gamma(t)$ is called a lightlike point if $\gamma'(t)$ is a lightlike vector.

If γ is spacelike or timelike, then it can be reparametrised by arc-length and its curvature is well defined at each point (see, for example, [11]). One can also have the notion of vertices (points where the derivative of the curvature vanishes). The curvature of γ is not defined at lightlike points (so we have no notion of vertices at such points). An inflection can be defined in terms of the contact of the curve with lines, so the curve can have an inflection at a lightlike points. For a generic curve, the lightlike points are not inflection points.

It is shown in ([11], Proposition 2.1) that the set of lightlike points of a closed curve γ is the union of at least four disjoint non-empty and closed subsets of γ (Figure 2). The complement of these sets are disjoint connected spacelike or timelike pieces of the curve γ .




Figure 2: Lightlike points in thick on a smooth closed curve in \mathbb{R}_1^2 .

The Minkowski set is defined in [11] as follows.

Definition 2.1 *The Minkowski Symmetry Set (MSS) of a curve γ in the Minkowski plane is the closure of the locus of centres of bi-tangent pseudo-circles to the curve.*

In \mathbb{R}_1^2 , we have to consider the fact that vectors can have negative length. For this reason, we say that a pseudo-circle is *maximal* if its radius equals the absolute minimum *modulus* distance from its centre to γ . The radius of such a circle cannot be increased if it is of type $S^1(p, r)$ or decreased if it is type $H^1(p, r)$ without it crossing γ .

Definition 2.2 *The Minkowski medial axis (MMA) of a curve γ in \mathbb{R}_1^2 is the subset of the Minkowski symmetry set formed by the centres of bi-tangent pseudo-circles which are maximal.*




Figure 3: A circle and its *MSS* (the two transverse line segments). The *MMA* is the subset of the *MSS* represented by a thick line.

Remark 2.3 In the Euclidean plane, for a closed curve γ , maximality implies that the bi-tangent circles are either entirely inside or entirely outside the curve ([7]). In the Minkowski plane however, since pseudo-circles are not compact, maximality ensures that the centres are entirely outside the curve, see Theorem 4.3.

The family of distance-squared functions $f : J \times \mathbb{R}_1^2 \rightarrow \mathbb{R}$ on γ is given by

$$f(t, c) = \langle \gamma(t) - c, \gamma(t) - c \rangle.$$

and the *extended family of distance-squared functions* $\tilde{f} : J \times \mathbb{R}_1^2 \times \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$\tilde{f}(t, c, r) = \langle \gamma(t) - c, \gamma(t) - c \rangle - r^2.$$

Denote by $f_c : J \rightarrow \mathbb{R}$ the function given by $f_c(t) = f(t, c)$. We say that f_c has an A_k -singularity at t_0 if $f'_c(t_0) = f''_c(t_0) = \dots = f_c^{(k)}(t_0) = 0$ and $f_c^{(k+1)}(t_0) \neq 0$. This is equivalent to the existence of a local re-parametrisation h of γ at t_0 such that $(f \circ h)(t) = \pm(t - t_0)^{k+1}$.

If f_c has a singularity at t_1 of type A_k and at t_2 of type A_l , we say that f has a multi-local singularity of type $A_k A_l$.

Geometrically, f_c has an A_k -singularity if and only if the curve γ has contact of order $k + 1$ at $\gamma(t_0)$ with the pseudo-circle $C(c, r)$ of centre c and radius r , with $|r| = \|\gamma(t_0) - c\|$. The distance squared function f_c has $A_k A_l$ -singularity if the pseudo circle $C(c, r)$ is tangent to γ at two distinct points and has order of contact $k + 1$ at one of them and $l + 1$ at the other.

It follows from Thom's transversality theorem (see for example [2, 8]) that for an open and dense set of immersions $\gamma : S^1 \rightarrow \mathbb{R}_1^2$ the function f_c has only local singularities of type A_1, A_2, A_3 and multi-local singularities of type $A_1^2, A_1 A_2, A_1^3$.

The *MSS* is the multi-local component of the bifurcation set of the family f , that is,

$$MSS = \{c \in \mathbb{R}_1^2 \mid \exists t_1, t_2 \text{ such that } t_1 \neq t_2, f_c(t_1) = f_c(t_2), f'_c(t_1) = f'_c(t_2) = 0\}.$$

It follows from Theorem 3.2 in [9] that the family f is always a versal unfolding of its generic singularities, so the *MSS* is locally diffeomorphic to the bifurcation set of the models of such singularities (Corollary 3.3 in [9]). Thus the configuration of the *MSS* at the generic multi-local singularities of f_c are as in in Figure 4.




Figure 4: Generic local models of the *MSS* in continuous line (the dashed curve is the caustic of the curve). Only the A_1^2, A_1^3 and A_3 singularities occur generically on the the *MMA*.

For a point on the *MSS* to also belong to the *MMA* the relevant bi-tangent pseudo-circle must be maximal. This means that the *MMA* forms a subset of the *MSS* and in particular this condition ensures that only A_1^2, A_3 and A_1^3 can belong to *MMA*. This is because for the other generic singularity types, namely A_2 and $A_1 A_2$, the pseudo-circle locally crosses the curve and therefore it cannot be maximal.

In [11] it was shown that the *MSS* is a regular curve at c_0 if and only if the bi-tangent pseudo-circle to γ at $\gamma(t_1)$ and $\gamma(t_2)$ is not osculating at $\gamma(t_1)$ or at $\gamma(t_2)$. If

this is the case, the tangent line to the MSS at p is the perpendicular bisector to the chord joining $\gamma(t_1)$ and $\gamma(t_2)$. (This is also true in the Euclidean case, see [5].)

3 Local reconstruction of the curve from the MSS

Suppose we are given the MSS (or MMA) of a spacelike or timelike smooth curve γ . Then the MSS is either a spacelike or a timelike curve [11]. If the MSS is not singular, we parametrise it by arc length $c(s) = (x(s), y(s))$ and denote by $r(s)$ the radius of the bi-tangent circle to γ centred at $c(s)$. Then it is possible to reconstruct local parametrisations γ_1 and γ_2 of the two corresponding arcs of γ as an envelope of the bi-tangent circles $C(c(s), r(s))$.

Proposition 3.1 *If the curve $c(s)$ is timelike and the bi-tangent pseudo-circle is of type $S^1(p, r)$ or if the curve $c(s)$ is spacelike and the bi-tangent pseudo-circle is of type $H^1(p, r)$, then the points of tangency are given by*

$$\gamma_i = c + \left(\frac{\partial r}{\partial s}\right) rT + (-1)^i \left(r\sqrt{\left(\frac{\partial r}{\partial s}\right)^2 + 1}\right) N, \quad i = 1, 2, \quad (1)$$

where T and N are the unit tangent and unit Minkowski normal to the MSS and r is the radius of the bi-tangent pseudo-circle, all evaluated at s .

If the curve $c(s)$ is spacelike and the bi-tangent pseudo-circle is of type $S^1(p, r)$ or if the curve $c(s)$ is timelike and the bi-tangent pseudo-circle is of type $H^1(p, r)$, then the points of tangency are given by

$$\gamma_i = c - r \left(\frac{\partial r}{\partial s}\right) T + (-1)^i \left(r\sqrt{\left(\frac{\partial r}{\partial s}\right)^2 - 1}\right) N, \quad i = 1, 2. \quad (2)$$

Proof Suppose that c is spacelike and the pseudo-circles are of type $H^1(p, r)$. Then the equation of the pseudo-circle of radius $r(s)$ centred at $c(s)$, is the set of points $w \in \mathbb{R}_1^2$ such that

$$F(s, w) = \langle c(s) - w, c(s) - w \rangle + r(s)^2 = 0.$$

The envelope of this family of these pseudo-circles is given by

$$D(F) = \{w \in \mathbb{R}^2 : \exists s \in \mathbb{R} \text{ such that } F(s, w) = \frac{\partial F}{\partial s}(s, w) = 0\}.$$

Differentiating F with respect to s and dropping the arguments yields

$$\frac{\partial F}{\partial s} = 2\langle c - w, T \rangle + 2r \frac{\partial r}{\partial s}.$$

Since γ is spacelike, $\langle T, T \rangle = 1$ and $\langle N, N \rangle = -1$. Writing $c - w = \lambda T + \mu N$, with $\lambda, \mu \in \mathbb{R}$, and substituting into $\frac{\partial F}{\partial s}$ we obtain $\lambda + r \frac{\partial r}{\partial s} = 0$ so that $\lambda = -r(\partial r / \partial s)$. Substituting into F yields $\lambda^2 - \mu^2 + r^2 = 0$ so that $\mu = \pm r \sqrt{(\partial r / \partial s)^2 + 1}$. It follows that the locii of the envelope points are as given (1). The same method can be applied to find the formula for the envelope in the remaining cases. \square

Remark 3.2 In the Euclidean case the rate of change of the radius function is restricted to be less than or equal to 1 in order for the envelope to be real, see [7]. In the Minkowski setting this restriction only applies when the pseudo-circles and the curve c are of opposite type (spacelike/timelike), otherwise any function r gives a real envelope.

Remark 3.3 Theorem 4.1 of [9] states that for any point p of a closed smooth curve γ there exists another point $q \in \gamma$ and a pseudo-circle that is tangent to γ at both p and q . From this, and the fact that it is possible to reconstruct the curve locally, it follows that it is possible to reconstruct any smooth closed curve from its Minkowski symmetry set.

4 The Minkowski medial axis

We now introduce the concept of the Minkowski hull.

Definition 4.1 *The Minkowski hull $MH(\gamma)$ of a closed curve γ in the Minkowski plane is the region of the plane such that for any point $p \in MH(\gamma)$ there exists a point $q \in \gamma$ such that the Minkowski distance between p and q is zero.*

Proposition 4.2 *The complement of the Minkowski hull for a closed smooth curve γ in the Minkowski plane consists of 4 disjoint open regions.*

Proof We consider a smooth closed curve $\gamma : S^1 \rightarrow \mathbb{R}_1^2$ and use coordinates such that the axes are parallel to the lightlike directions. We write $\gamma(t) = (x(t), y(t))$. Since the functions $x(t)$ and $y(t)$ are bounded, they must both attain an absolute maximum and an absolute minimum. Since the curve is smooth, this gives exactly four extrema points (two of $x(t)$ and two of $y(t)$) on the curve. The curve γ is now contained the compact region determined by the tangent lines to these four extremal. The tangent lines divide the plane into the Minkowski hull and four disjoint regions of its complement, see Figure 5 (right). \square

Lemma 4.3 *The points on the MMA of a closed plane curve which are the centres of bi-tangent circles of type $H^1(p, r)$ or $S^1(p, r)$ are all inside the closure of the complement of the Minkowski hull of the curve.*




Figure 5: Left: the Minkowski hull of a closed curve in the Minkowski plane and its complement (shown in grey). Right: a circle, its Minkowski caustic (continuous curve with four cusps), Minkowski hull (shaded in gray), MSS and MMA (in thick).

Proof From the definition, a point p belonging to the Minkowski hull has Minkowski distance zero to some point on the curve. It follows that pseudo-circles of types $H^1(p, r)$ and $S^1(p, r)$ cannot be maximal because the light cone $LC^*(p)$ intersects the curve. Therefore, pseudo-circles of types $H^1(p, r)$ and $S^1(p, r)$ corresponding to the MMA lie in the complement of the Minkowski hull. \square

Proposition 4.4 *The MMA of a closed smooth convex plane curve lies strictly inside the closure of the complement of the Minkowski hull.*

Proof Closed smooth convex plane curves have exactly four closed lightlike regions (see [11]). Since the four centres of the pseudo-circles that are bi-tangent to these four regions form the boundary to the Minkowski hull, the centres also lie on the boundary. This, together with Lemma 4.3, prove that the centres all three types of maximal bitangent pseudo-circles lie either inside the complement of the Minkowski hull or on its boundary. Therefore, the MMA of a closed smooth convex plane curve lies strictly inside the closure of the complement of the Minkowski hull. See Figure 5 (right). \square

Corollary 4.5 *of 4.3. The MMA of a closed smooth (not necessarily convex) plane curve lies strictly inside the closure of the complement of the Minkowski hull of the curve except for components formed by the centres of bi-tangent pseudo-circles of type $LC(p)$ (note that these components are subsets of lightlike lines).*

Definition 4.6 *A pseudo-circle of type $H^1(p, r)$ or $S^1(p, r)$ is said to be 1-branch bi-tangent to a curve γ if one of its branches is tangent to γ in at least two distinct points.*

Since the *MMA* lies inside the closure of the complement of the Minkowski hull we have the following result.

Theorem 4.7 *The pseudo-circles of types $H^1(p, r)$ and $S^1(p, r)$ corresponding to the *MMA* of a closed plane curve are all 1-branch bitangent.*

Proof Lemma 4.3 states that the centres of the bitangent pseudo-circles of type $H^1(p, r)$ and $S^1(p, r)$ occur in the closure of the complement of the Minkowski hull of the curve. Since the bitangent pseudo-circle necessarily has one of its branches completely contained in the complement of the Minkowski hull it follows that the tangencies must occur on only one of the pseudo-circle's two branches. See Figure 5. \square

Remark 4.8 Note that the converse of Theorem 4.7 is true for convex curves but is not true for non-convex curves. That is, not all 1-branch bitangent pseudo-circles are maximal for non-convex curves. See for example Figure 7 (right).

One of the useful properties of the Euclidean medial axis is that it can be used to reconstruct the original curve. It is shown in Proposition 4.5 of [9] that for the piece of curve $\alpha(t) = (t, t^3)$, $-\frac{1}{2} < t < \frac{1}{2}$, there are no 1-branch bitangent pseudo-circles. The curve $\alpha(t)$ can be extended smoothly to obtain a closed curve γ . It is possible to construct γ so that there do not exist pseudo-circles which are 1-branch tangent to a point of $\alpha(t)$ and are also tangent to some other point of $\gamma(t)$. Consider for example the limaçon whose radius r is given by $r = \frac{3}{2} + \cos(\theta)$ where $-\pi < \theta < \pi$. Splitting the limaçon into timelike and spacelike components, only pairs of points from the regions $\arctan(\frac{\sqrt{35}}{17}) - \pi < \theta < -\arctan(\frac{\sqrt{35}}{17}) + \pi$ have corresponding Minkowski medial axis points, see Figure 6.

Therefore, the *MMA* together with a radius function (*MMA* transform), unlike its Euclidean counterpart is not a complete shape describer.

The fact that the *MMA* consists of centres of only 1-branch bi-tangent pseudo-circles motivates the following new type of medial axis construction, called the *1-branch Minkowski medial axis*.

Definition 4.9 *The 1-branch Minkowski medial axis is the locus of the centres of pseudo-circles that are tangent to the curve γ in two or more points such that the tangencies occur on just one of the branches of the pseudo-circle.*

Since the *MMA* is made up of *only* 1-branch bitangent pseudo-circle centres, it is a subset of the 1-branch medial axis.

The two sets are not equal because not all 1-branch bitangent pseudo-circles are maximal, see for example Figure 7 (left). The 1-branch Minkowski medial axis does not lie inside the complement of the Minkowski hull for non-convex curves (see for example Figure 7 (right)).




Figure 6: The limaçon given by $r = \frac{3}{2} + \cos(\theta)$ where $-\pi < \theta < \pi$, its Minkowski caustic and Minkowski medial axis. Only pairs of points in the region $\arctan(\frac{\sqrt{35}}{17} - \pi < \theta < -\arctan(\frac{\sqrt{35}}{17}) + \pi)$ have corresponding Minkowski medial axis points.



Figure 7: Left: an illustration of a 1-branch bi-tangent pseudo circle which is not maximal (A_1A_2 -singularity). Right: a 1-branch bi-tangent pseudo-circle whose centre lies inside a non-convex curve.

Remark 4.10 Theorem 4.5 of [9] states that for any point p on a spacelike or timelike curve γ without inflections there exists another point q on γ and a pseudo-circle that is tangent to γ at both p and q with both points being on a single branch of the pseudo-circle. From this it follows that any closed convex curve can be reconstructed from its 1-branch medial axis. The 4 lightlike components are either isolated points or lightlike line segments (in the generic case only isolated points are possible). To complete γ , these components can be added by taking the closure of the curve if they are just isolated points, or by joining up the remaining components with lightlike lines.

5 Shocks on the Minkowski medial axis

At each point on the medial axis of a curve γ in the Euclidean plane there is an associated radius function r corresponding to the radius of the bi-tangent circle. The direction of the increasing radius function on the medial axis, that is the direction for which $\frac{\partial r}{\partial s}(s) > 0$, can be indicated by an arrow and this gives the shock set (see for example [1, 6]).

The shock set is a dynamic view of the medial axis. If there is a propagation of waves (grass fire) from the curve γ , then this leads to the formation of singularities (the medial axis). The shock set gives the direction along which this formation of singularities propagates.

In [6] the local generic forms of shocks that can occur on the medial axis in the Euclidean plane are classified. It is shown that some types of shocks cannot occur generically on the Euclidean medial axis. For example, it is proven that the only form of shock that can occur at an A_3 -singularity of a given distance squared function on γ is that with outward velocity, see Figure 8.

We define the analogue of shocks in the Minkowski plane. This is the Minkowski medial axis together with an arrow in the direction of increasing radius of the corresponding bitangent pseudo-circle.

In this section we show that for the *MMA* both types of shock can occur (outwards and inwards) at an A_3 -singularity of a given distance squared function on γ , depending on whether the *MMA* is spacelike or timelike. We also show that the generic shocks that can occur at an A_1^3 -singularity of a given distance squared function on γ are different to those on the Euclidean medial axis.

In what follows, the singularities refer to those of a given distance squared function.

5.1 Shocks at an A_3 -singularity

The A_3 -singularity occurs at a vertex of the curve. These occur where the two A_1 -contact points for nearby A_1^2 -bi-tangent pseudo-circles come into coincidence. The A_1 -points must therefore lie on the same branch of the pseudo-circle. This means that given a *MMA* near an A_3 point and its associated radius function, formula (2) can be used to find the corresponding points on γ .

Theorem 5.1 *If the curve γ is timelike (resp. spacelike) at a vertex, then the shock on the *MMA* is of outwards (resp. inwards) type.*

Proof Consider a neighbourhood of an A_3 point on a spacelike *MMA* (which necessarily corresponds to a timelike piece of γ , see Theorem 5.2 in [9]). Orient the *MMA* so that its tangent line points towards the branch of the bi-tangent pseudo-circle that contains the tangent points. Since the envelope points of the bi-tangent pseudo-circles are in the direction of the tangent line it follows from formula (2) that $-r\frac{\partial r}{\partial s} > 0$. As




Figure 8: Comparing A_3 shocks that can occur on the Euclidean and Minkowski medial axes. In the Minkowski plane, both cases can occur and are distinguished by the type of the bi-tangent pseudo-circle.

$r > 0$, $\frac{\partial r}{\partial s} < 0$ so the radius function r must decrease in the direction of the A_3 point. Therefore, for a timelike γ near a vertex, the shocks are of outwards type; see Figure 8.

If the MMA is timelike (the curve γ must be spacelike, Theorem 5.2 in [9]). Following the same arguments as above and using formula (2) at an A_3 -point, we get $-r \frac{\partial r}{\partial s} > 0$. Here $r < 0$ so it follows $\frac{\partial r}{\partial s} > 0$. This implies that the shocks are of inwards type; see Figure 8. \square

Remark 5.2 *Theorem 5.1 also holds when the MMA is replaced by the 1-branch MMA .*

5.2 Shocks at an A_1^3 -singularity

As with the shocks at A_3 -singularity, the shocks at A_1^3 -singularity turn out to be different from those of the Euclidean medial axis. The type of shocks that can occur depends on whether the bi-tangent pseudo-circle is of type $H^1(p, r)$ or $S^1(p, r)$.

For closed curves, the MMA only consists of the centres of pseudo-circles whose tangencies occur on only one of its branches (Theorem 4.7). Theorems 5.4 and 5.5 give a classification of shocks that can occur for closed curves. For two disjoint pieces of curves, it is possible that the pseudo-circle centred on the medial axis can be tangent to each piece of curve (so the centre is not on the 1-branch MMA). For completeness, Theorem 5.7 gives the classification of shocks at an A_1^3 -singularity when the relevant pseudo-circle has at least one tangency on each of its branches.




Figure 9: A_1^3 shocks that can occur on the Euclidean and Minkowski medial axes for closed curves. Case (2) occurs when the pseudo-circle is of type S^1 (here the tri-tangency occurs on the branch above the figure). Case (3) occurs when the pseudo-circle is of type H^1 (here the tri-tangency occurs on the branch to the left of the figure)

Suppose that γ is a closed plane curve and the three tri-tangent points, say q_1, q_2 and q_3 on γ to a pseudo-circle all lie on one branch of the pseudo-circle. Each branch of the MMA corresponds to the centres of bi-tangent pseudo-circles whose points of tangency are near two of the points $q_i, i = 1, 2, 3$. For each branch, the two corresponding tangency points are called the characteristic points and are denoted X^- and X^+ . If these points are near q_i and q_j , we denote by P the point q_k , with $k \neq i, j$.

Lemma 5.3 *If the arc that contains X^- and X^+ does not contain (resp. contains) the point P , then the medial axis goes in the direction of entering (resp. goes away from) the arc.*

Proof We consider the case of a tri-tangent pseudo-circle of type $S^1(p, r)$, the proof is similar for a tri-tangent pseudo-circle of type $H^1(p, r)$ and is omitted.

Consider the function $f(s) = \langle c(s) - P, c(s) - P \rangle - (r(s))^2$, where s is the arc-length parameter of the MSS branch $c(s)$ and $r(s)$ is the radius of the bi-tangent pseudo-circle. Let s_0 correspond to the A_1^3 point. Note that $f(s_0) = 0$. We have tangency of type A_1 , so $f'(s_0) \neq 0$. If $f'(s_0) < 0$, then $f(s) < f(s_0)$ for small $s > s_0$. For such s , $c(s)$ cannot be on the MMA since the point P will have come ‘inside’ the pseudo-circle centre $c(s)$ radius $r(s)$. Here, ‘inside’ means that its absolute distance from the centre is less than $|r|$.

Now $f'(s) = 2\langle c(s) - P, T(s) \rangle - 2r(s)r'(s)$, so that $f'(s_0) < 0$ is equivalent to $\langle c(s_0) - P, T(s_0) \rangle < r(s_0)r'(s_0)$. We have $r(s_0)r'(s_0) = \langle c(s_0) - X^\pm, T(s_0) \rangle$. This in turn implies $\langle X^\pm - P, T \rangle < 0$.

We parametrise the tritangent pseudo-circle by $g(t) = (r_0 \sinh(t), r_0 \cosh(t))$. Suppose that $X^+ = g(\theta)$, $X^- = g(\varphi)$ and $P = g(\rho)$ for some θ, φ and ρ .

The tangent to the branch of the *MSS* corresponding to the characteristic points X^+ and X^- has direction $T = (\sinh(\frac{\theta+\varphi}{2}), \cosh(\frac{\theta+\varphi}{2}))$ and the vector $(X^+ - p)$ has direction $(\cosh(\frac{\theta+\rho}{2}), \sinh(\frac{\theta+\rho}{2}))$ if $\theta > \rho$ and $(-\cosh(\frac{\theta+\rho}{2}), -\sinh(\frac{\theta+\rho}{2}))$ if $\theta < \rho$. Assume that $\theta > \rho$. The Minkowski product $\langle X^\pm - P, T \rangle$, up to a positive factor, is given by

$$-\sinh\left(\frac{\theta+\varphi}{2}\right)\cosh\left(\frac{\theta+\rho}{2}\right) + \sinh\left(\frac{\theta+\rho}{2}\right)\cosh\left(\frac{\theta+\varphi}{2}\right) = 2(e^\rho - e^\varphi)e^{-(\theta+\varphi+\rho)}$$

which is negative if and only if $\rho < \varphi$.

Similarly, assuming $\theta < \rho$ gives that $\langle X^\pm - P, T \rangle$ is negative if and only if $\varphi < \rho$.

Thus the condition $\langle X^\pm - P, T \rangle < 0$ is equivalent to either $\rho < \varphi, \theta$ or $\rho > \varphi, \theta$ which is equivalent to the statement in Lemma 5.3. \square

Proposition 5.4 *For a closed curve, the Minkowski medial axis at an A_1^3 -singularity has shock type (2) in Figure 9 if the tri-tangent pseudo-circle is of type $S^1(p, r)$.*

Proof For three points on the same branch of the pseudo-circle and for ρ, φ and θ as in the proof of Lemma 5.3, the conditions $\rho < \varphi, \theta$ or $\rho > \varphi, \theta$ must be true for two of the three medial axis branches. The arrows, indicating directions of increasing radius, can now be added to the medial axes. Formula (2) implies that the radius must be increasing in the direction of the branch of the pseudo-circle that contains the three tritangent points. \square

Similarly the following proposition holds:

Proposition 5.5 *For a closed curve, the Minkowski medial axis at an A_1^3 point has shock type (3) in Figure 9 if the tri-tangent pseudo-circle is of type H^1 .*

Remark 5.6 Propositions 5.4 and 5.5 also hold when the *MMA* is replaced by the 1-branch *MMA*. In this case the condition that the curve be closed can also be dropped.

Proposition 5.4 and Proposition 5.5 give a complete classification of shocks for closed curves at an A_1^3 -singularity.

Observe that the condition to be on the *MSS* is dependent on two or more points on the boundary, in contrast the condition for a point to be on the *MMA* is global in that it depends on all of the points on the boundary. Also note that both the

MSS and the MMA are well defined for both closed curves and open curves (ignoring end-points).

Due to the fact that pseudo-circles consist of two components, the classification of shocks on the MMA is slightly different from that of Euclidean medial axis [6]. For n disjoint tangent lines that share a common bitangent circle it is always possible to construct a *closed* curve that shares these tangent lines such that the circle is maximal (contained inside the curve). If they share a common pseudo-circle however, it is always possible to construct a *closed* curve that shares these tangent lines such that the pseudo-circle is maximal if and only if they are tangent to just one of the branches. However, if they are tangent to both branches it is possible to construct a curve consisting of *two* branches that shares these tangent lines such that the pseudo circle is maximal.

So using the local type construction as used in Lemma 5.3 to determine the shocks in the Euclidean case there is only one case which is valid for open and closed curves, see for example [6]. In the Minkowski case however, the closed and two branched curve cases need to be considered separately.

Blum viewed the (Euclidean) medial axis as a quench point for grass-fire flow initiated from the boundary of the shape [1]. He considered the medial axes of curve segments as well as closed curves. In this spirit, and for the purpose of applications, we now consider the MMA and classify its shocks for when the tri-tangency occurs on both branches of the pseudo-circle.

Theorem 5.7 *The shocks that can occur at an A_1^3 -singularity when the tangent points occur on both branches of a pseudo-circle of type $S^1(p, r)$ are as shown in Figure 10.*

Remark 5.8 *For pseudo-circles of type $H^1(p, r)$ the proof works the same and the shocks are the same as in Figure 10 with the directions of the arrows reversed.*

Proof of Theorem 5.7. We take the centre of the pseudo-circle to be the origin and parametrise its branches by $g_1(t) = (r_0 \sinh(t), r_0 \cosh(t))$ and $g_2(t) = (r_0 \sinh(t), -r_0 \cosh(t))$ and assume that the branch containing two tangent points to be the branch in the lower half of the plane. Denote by X_θ the tangent point on the upper branch at the point $g_1(\theta)$ for some value θ , and X_φ and X_ρ the two tangent points on the lower branch at points $g_2(\varphi)$ and $g_2(\rho)$, respectively.

Denote by $T_{\theta, \varphi}$ the tangent line to the MMA branch that corresponds to the characteristic points θ and φ and similarly for $T_{\theta, \rho}$ and $T_{\rho, \varphi}$. Consider first the branch of the MMA with tangent line $T_{\rho, \varphi}$. This tangent line has direction $T_{\rho, \varphi} = (\sinh(\frac{\rho+\varphi}{2}), \cosh(\frac{\rho+\varphi}{2}))$ and the vector joining the point X_ρ to X_θ , is given by $(X_\varphi - X_\theta) = r_0(\sinh(\varphi) - \sinh(\theta), \cosh(\varphi) + \cosh(\theta))$. Taking their Minkowski product yields

$$\langle (X_\varphi - X_\theta), T_{\rho, \varphi} \rangle = \langle (X_\rho - X_\theta), T_{\rho, \varphi} \rangle = r_0 \cosh\left(\frac{\varphi - \rho}{2}\right) + r_0 \cosh\left(\frac{\varphi + \rho}{2} + \theta\right)$$

which is always positive. Therefore, the corresponding branch of the *MMA* starts at the A_1^3 point and goes in the direction of the branch containing the characteristic points. Formula (2) implies that the shock travels along this *MMA* branch in the direction towards the centre.

Consider now the two branches with tangents $T_{\theta,\rho}$ and $T_{\theta,\varphi}$. For each branch of the *MMA* there are four possibilities: The *MMA* can be to the left or to the right of the centre (since the branches are both timelike), and in both cases the direction of the shock can be either towards or away from the centre.

The tangent line $T_{\theta,\varphi}$ to the medial axis at the A_1^3 , up to a nonzero factor, can be written $T_{\theta,\varphi} = \left(\frac{\sinh(\theta)+\sinh(\varphi)}{2}, \frac{\cosh(\theta)-\cosh(\varphi)}{2} \right)$ and the vector joining the point X_ρ to X_θ , is given by $(X_\theta - X_\rho) = r_0(\sinh(\theta) - \sinh(\rho), \cosh(\theta) + \cosh(\rho))$. Taking their Minkowski product yields:

$$\langle (X_\theta - X_\rho), T_{\theta,\varphi} \rangle = \frac{r_0}{2}(1 - \cosh(\rho - \theta) + \cosh(\rho + \varphi) - \cosh(\theta + \varphi)) \quad (3)$$

which is positive when $-\varphi < \theta < \rho$ or $\rho < \theta < -\varphi$ and negative otherwise. Recall that when this product is positive small positive $s > s_0$ belongs to the *MMA* and when negative, and when negative it is small $s < s_0$ that belong to the *MMA*. Now, the direction of increasing radius can be added to the MMS. It follows directly from formula (1) that if $\theta + \varphi > 0$ the radius function on the branch corresponding to $T_{\theta,\varphi}$ increases from left to right, whereas if $\theta + \varphi < 0$ the radius increases from right to left, and similarly for $T_{\theta,\rho}$.

Considering the two branches together now, it must be determined which branch goes ‘over’ the other. Comparing the gradients of the two tangent lines it can be seen that $T_{\theta,\varphi}$ is steeper than $T_{\theta,\rho}$ if and only if

$$\theta < \rho < \varphi, \quad \rho < \varphi < \theta \quad \text{or} \quad \varphi < \theta < \rho.$$

Otherwise, $T_{\theta,\rho}$ is the steeper of the two.

Considering these conditions, along with the above conditions for which side the branches lie on, gives the complete list of possible shocks that can occur (see table 1 and Figure 10). Note that for spacelike A_1^3 singularities, the shocks can be obtained from rotating those in Figure 10 by $\frac{\pi}{2}$ and reversing the directions of the arrows. \square

References

- [1] H. Blum, A transformation for extracting new descriptors of shape. *W. Whaten-Dunn (Ed.), Models for the Perception of Speech and Visual Forms*. MIT Press, Cambridge, MA, 1967, 362-380.




Figure 10: The types shocks that can occur for timelike A_1^3 points.

- [2] J. W. Bruce and P. J. Giblin, *Curves and Singularities*. Cambridge University Press, 1992.
- [3] I. A. Bogaevsky, Perestroikas of shocks and singularities of minimum functions. *Physica D.*, 173, (2002) 1-29.
- [4] J. W. Bruce, P. J. Giblin and C. G. Gibson, Symmetry Sets. *Proc. Roy. Soc. Edinburgh*, 101, (1985) 163-186.
- [5] P. J. Giblin and S. Brassett, Local symmetries of plane curves. *Amer. Math. Monthly* 92, (1985) 698-707.
- [6] P. J. Giblin and B .B. Kimia, On the local form and transitions of symmetry sets, medial axes, and shocks. *Int. J. of Comput. Vis.* 54, (2003) 143-157.
- [7] P. J. Giblin and J. P. Warder, Reconstructions from Medial Representations. *Am. Math. Monthly*, 118, (2011) 712-725.

Table 1: Conditions for shocks occurring on the MMA at a timelike A_1^3 singularity

a)	$-\rho < -\varphi < \theta < \varphi < \rho$ $-\varphi < -\rho < \theta < \rho < \varphi$	b)	$-\rho < \theta < \varphi < -\varphi < \rho$ $-\rho < \theta < -\varphi < \varphi < \rho$ $-\varphi < \theta < -\rho < \rho < \varphi$ $-\varphi < \theta < -\rho < \rho < \varphi$
c)	$\theta < -\varphi, \varphi, -\rho, \rho$	d)	$-\varphi, \varphi, -\rho, \rho < \theta$
e)	$\varphi < -\rho < \rho < \theta < -\varphi$ $\varphi < \rho < -\rho < \theta < -\varphi$ $\rho < -\varphi < \varphi < \theta < -\rho$ $\rho < \varphi < -\varphi < \theta < -\rho$	f)	$\rho < \varphi < \theta < -\varphi < -\rho$ $\varphi < \rho < \theta < -\rho < -\varphi$
g)	$-\varphi < -\rho < \rho < \theta < \varphi$ $-\varphi < \rho < -\rho < \theta < \varphi$ $-\rho < -\varphi < \varphi < \theta < \rho$ $-\rho < \varphi < -\varphi < \theta < \rho$	h)	$-\varphi < \rho < \theta < -\rho < \varphi$ $\rho < -\varphi < \theta < \varphi < -\rho$ $-\rho < \varphi < \theta < -\varphi < \rho$ $\varphi < -\rho < \theta < \rho < -\varphi$
i)	$\rho < \theta < \varphi < -\varphi < -\rho$ $\rho < \theta < -\varphi < \varphi < -\rho$ $\varphi < \theta < -\rho < \rho < -\varphi$ $\varphi < \theta < -\rho < \rho < -\varphi$		

- [8] M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*. Graduate Texts in Mathematics, Vol. 14. Springer-Verlag, New York-Heidelberg, 1973.
- [9] G. M. Reeve and F. Tari, Mikowski symmetry sets of plane curves. Preprint, 2013.
- [10] K. Siddiqi and S. Pizer, *Medial Representations: Mathematics, Algorithms and Applications*. Springer: Computational Imaging and Vision 37, 2008.
- [11] A. Saloom and F. Tari, Curves in the Minkowski plane and their contact with pseudo-circles. *Geom. Dedicata* 159, (2012) 109-124.

Instituto de Ciências Matemáticas e de Computação - USP, Avenida Trabalhador são-carlense, 400 - Centro, CEP: 13566-590 - São Carlos - SP, Brazil.
Emails: grahamreeve86@gmail.com, faridtari@icmc.usp.br